



---

## PHYSICS

Paper 1 Multiple Choice

**9749/01**

**20 September 2018**

**1 hour**

Additional Materials: Multiple Choice Answer Sheet

---

### READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use paper clips, glue or correction fluid.

Write your name, civics group and registration number on the Answer Sheet in the spaces provided.

There are **thirty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

**Read the instructions on the Answer Sheet very carefully.**

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

The use of an approved scientific calculator is expected, where appropriate.

---

This document consists of **18** printed pages and **0** blank page.

**Data**

speed of light in free space,	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
permeability of free space,	$\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$
permittivity of free space,	$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$ $(1/(36 \pi)) \times 10^{-9} \text{ F m}^{-1}$
elementary charge,	$e = 1.60 \times 10^{-19} \text{ C}$
the Planck constant,	$h = 6.63 \times 10^{-34} \text{ J s}$
unified atomic mass constant,	$u = 1.66 \times 10^{-27} \text{ kg}$
rest mass of electron,	$m_e = 9.11 \times 10^{-31} \text{ kg}$
rest mass of proton,	$m_p = 1.67 \times 10^{-27} \text{ kg}$
molar gas constant,	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
the Avogadro constant,	$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
the Boltzmann constant,	$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$
gravitational constant,	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
acceleration of free fall,	$g = 9.81 \text{ m s}^{-2}$

**Formulae**


uniformly accelerated motion,	$s = ut + \frac{1}{2}at^2$
	$v^2 = u^2 + 2as$
work done on/by a gas,	$W = p\Delta V$
hydrostatic pressure,	$p = \rho gh$
gravitational potential,	$\phi = -\frac{Gm}{r}$
temperature,	$T/K = T / ^\circ\text{C} + 273.15$
pressure of an ideal gas,	$p = \frac{1}{3} \frac{Nm}{V} \langle c^2 \rangle$
mean translational kinetic energy of an ideal gas molecule	$E = \frac{3}{2}kT$
displacement of particle in s.h.m.	$x = x_0 \sin \omega t$
velocity of particle in s.h.m.	$v = v_0 \cos \omega t$ $= \pm \omega \sqrt{(x_0^2 - x^2)}$
electric current,	$I = Anvq$
resistors in series,	$R = R_1 + R_2 + \dots$
resistors in parallel,	$1/R = 1/R_1 + 1/R_2 + \dots$
electric potential,	$V = \frac{Q}{4\pi\epsilon_0 r}$
alternating current/voltage,	$x = x_0 \sin \omega t$
magnetic flux density due to a long straight wire	$B = \frac{\mu_0 I}{2\pi d}$
magnetic flux density due to a flat circular coil	$B = \frac{\mu_0 NI}{2r}$
magnetic flux density due to a long solenoid	$B = \mu_0 nI$
radioactive decay,	$x = x_0 \exp(-\lambda t)$
decay constant	$\lambda = \frac{\ln 2}{t_{\frac{1}{2}}}$

- 1 Measurements are subject to both systematic error and random error.

Which measurements have high accuracy and low precision?




- A High random error and high systematic error
- B High random error and low systematic error
- C Low random error and high systematic error
- D Low random error and low systematic error

- 2 The diagram below shows the variation of displacement with time for 2 trains, Train A and Train B, running on parallel tracks.




Which of the following statements is correct?

- A At time  $t_B$ , both trains have the same velocity.
  - B Both trains speed up all the time.
  - C Both trains have the same velocity at some time before  $t_B$ .
  - D Somewhere on the graph, both trains have the same acceleration.
- 3 A ball thrown at an angle travels in a trajectory as shown below. When the ball is at the top of its flight, which of the following shows the direction its resultant acceleration?




- 4 The diagram shows a man standing on a platform that is attached to a flexible pipe. Water is pumped through the pipe so that the man and platform remain at a constant height.



The resultant vertical force on the platform is zero. The combined mass of the man and platform is 96 kg. The mass of water that is discharged vertically downwards from the platform each second is 40 kg.

What is the speed of the water leaving the platform?


- A** 2.4 m s<sup>-1</sup>      **B** 6.9 m s<sup>-1</sup>      **C** 24 m s<sup>-1</sup>      **D** 47 m s<sup>-1</sup>
- 5 Two objects A and B collide head-on on a horizontal, frictionless surface. The velocities of the objects before the collision and after the collision are shown in the diagram below.



Which of the following statements is correct?


- A** The sum of the kinetic energies of A and B is conserved.  
**B** The collision is possibly inelastic.  
**C** The total momentum of the two objects cannot be conserved since no net external force acts on them.  
**D** The momentum of each object is conserved.

- 6 A wooden block of mass  $1.0 \text{ kg}$  is on a rough horizontal surface. A force of  $12 \text{ N}$  is applied to the block and it accelerates at  $4.0 \text{ m s}^{-2}$ .




What is percentage of work done is used to overcome frictional force when the block is being moved a distance of  $10 \text{ m}$ ?

- A 33.3%      B 66.7%      C 75.0%      D 100%
- 7 A small bead is set into circular motion in a horizontal plane within a smooth conical cone as shown below.



If the bead is moving with a centripetal acceleration of  $g$ , what is the angle  $\theta$ ?

- A  $30^\circ$       B  $45^\circ$       C  $55^\circ$       D  $70^\circ$
- 8 In the diagram below, the volume of bulb X is twice that of bulb Y. The system is filled with an ideal gas and a steady state is established with the bulbs held at  $200 \text{ K}$  and  $400 \text{ K}$ .




There are  $b$  moles of gas in X.

How many moles of gas are there in Y?

- A  $b/4$       B  $b/2$       C  $b$       D  $2b$

- 9 The diagram below shows a motorised vehicle for carrying one person.



The vehicle has two wheels on one axle. The passenger stands on a platform between the wheels.

The weight of the machine is 600 N. Its centre of mass is 200 mm in front of the axle. The wheel radius is 400 mm.


When stationary, a passenger of weight 600 N stands with his centre of mass 200 mm behind the axle to balance the machine.

The motor is now switched on to provide a horizontal force of 90 N at the ground to move the vehicle forwards.

How far and in which direction must the passenger move his centre of mass to maintain balance?

- A 60 mm backward
- B 60 mm forward
- C 140 mm backward
- D 140 mm forward

- 10 The gravitational potential  $\phi$  as it varies with distance  $r$  from the centre of the Moon to the centre of Earth as shown in the diagram below.



Which of the following statements is **not** true?

- A The gravitational field strength at  $P$  is zero.
  - B The gravitational field strength at  $P$  is the scalar sum of the individual field strengths of the Moon and the Earth at point  $P$ .
  - C The gravitational field strength at  $P$  is the vector sum of the individual field strengths of the Moon and the Earth at point  $P$ .
  - D The gravitational field strength at  $P$  is given by the rate of change of  $\phi$  with respect to  $r$ .
- 11 A fixed amount of ideal gas undergoes the following changes:

Process 1: The gas is heated at constant volume.

Process 2: The gas is compressed at constant pressure.





How does the internal energy of the gas for each process change?

	Process 1	Process 2
A	increase	increase
B	increase	decrease
C	decrease	increase
D	decrease	decrease


- 12 Which of the following statements is **false** in describing a mass moving in simple harmonic motion?
- A The maximum kinetic energy is dependent on the frequency of the oscillation.
  - B The time taken for the system to change from maximum kinetic energy to maximum potential energy is a quarter of the period of the oscillation.
  - C An oscillating system with larger amplitude will have a longer period.
  - D An oscillating system with larger amplitude will have a greater maximum velocity.



- 13 Which of the following graphs show the variation in the total energy of an object under light damping as time passes?




- 14 A 500 Hz tuning fork is held at the open end of an air-filled glass tube, which is closed at the other end by a movable piston.







Resonance is achieved when  $L = 82.5$  cm. Given that the speed of sound in air is  $330 \text{ m s}^{-1}$ , which of the following other values of  $L$  will **not** result in resonance?

- A** 16.5 cm      **B** 49.5 cm      **C** 99.0 cm      **D** 115.5 cm


- 15 Unpolarised light is incident normally on a linear polarizer  $P$  and subsequently on a second linear polarizer  $Q$  as shown below.



Which graph best represents the relationship between the emergent light intensity  $I_{\text{emerge}}$  and the angle of rotation of  $Q$  about the axis of light transmission  $\theta$ ?




- 16 In the diagram below (drawn to scale), X and Y are identical point sources of waves that exhibit a constant phase difference of  $\pi$  at source. The waves have a wavelength of 1.0 cm.



Which of the following statements is true?


- A P is a point of destructive interference.  
 B  $XQ - YQ = m \frac{\lambda}{2}$ , where  $m$  is an odd integer.  
 C  $XR - YR = n\lambda$ , where  $n$  is an integer.  
 D S is a point of destructive interference.
- 17 The diagram below shows electric field lines with points P, Q and R on one of the field lines. The distance between PQ is equal to the distance between QR.



If the potential at P and Q are 0 V and  $-200$  V, respectively, which of the following can be a possible value of the potential at R?

- A  $-450$  V      B  $-400$  V      C  $+200$  V      D  $+250$  V


18 Four identical point charges are arranged at the corners of a square of length  $R$  as shown below.



What is the magnitude of the electric field strength  $E$  and the electric potential  $V$  at point  $X$ ?

	$E$	$V$
<b>A</b>	$\frac{12}{5} \frac{Q}{\pi\epsilon_0 R^2}$	$\frac{2}{5} \frac{Q}{\pi\epsilon_0 R}$
<b>B</b>	$\frac{12}{5} \frac{Q}{\pi\epsilon_0 R^2}$	$\left(1 + \frac{1}{\sqrt{5}}\right) \frac{Q}{\pi\epsilon_0 R}$
<b>C</b>	$\frac{4}{\sqrt{5}^3} \frac{Q}{\pi\epsilon_0 R^2}$	$\frac{2}{5} \frac{Q}{\pi\epsilon_0 R}$
<b>D</b>	$\frac{4}{\sqrt{5}^3} \frac{Q}{\pi\epsilon_0 R^2}$	$\left(1 + \frac{1}{\sqrt{5}}\right) \frac{Q}{\pi\epsilon_0 R}$


19 A copper wire has a number density of  $8.5 \times 10^{28}$  conduction electrons per cubic metre, and an cross-sectional area of  $3.2 \times 10^{-7} \text{ m}^2$ . When a potential difference is applied to the ends of the wire, the current is 1.0 A.



If all the electrons within a cylinder of length  $L$  pass point  $X$  in 60 s, what is the value of  $L$ ?


- A** 0.00015 m      **B** 0.014 m      **C** 0.025 m      **D** 0.20 m

20 Which of the following statements is true about the circuit shown?







- A When switch  $K$  is closed, as the resistance  $R_2$  increases, the balanced length  $L$  decreases.
- B When switch  $K$  is open, as the resistance  $R_2$  increases, the balanced length  $L$  does not change.
- C When switch  $K$  is open, as resistance  $R_1$  increases, the balanced length  $L$  decreases.
- D When switch  $K$  is closed, changes in the internal resistance of  $E_2$  will produce an increase in the balanced length  $L$ .


- 21 In a cathode-ray oscilloscope tube, the electron beam passes through a region where there is an electric field directed vertically downwards and a magnetic field directed vertically upwards as shown in the diagram below.



Which of the diagrams below shows a possible position of the spot on the screen when both fields are operating together?




- 22 A wooden cylinder of mass 0.250 kg and length  $L$  of 0.100 m, has 10 turns of wire wrapped around it longitudinally. The cylinder is released on a plane inclined at an angle  $\theta$  to the horizontal, with the plane of the coil parallel to the incline plane, as shown in the diagram below.



If there is a vertical uniform magnetic field of magnitude 0.500 T acting throughout the plane, what is the least current through the coil that keeps the cylinder from rolling down the plane?


- A 1.25 A      B 2.45 A      C 6.77 A      D 11.2 A
- 23 A closed circular loop of wire has a radius of 3.7 cm. It is bent along a diameter such that the two halves are perpendicular to each other. A uniform magnetic flux density of  $B = 76 \text{ mT}$  is directed perpendicular to the fold diameter and makes equal angles ( $45^\circ$ ) with the planes of the semicircle.



If the magnetic flux density  $B$  is reduced to zero at a uniform rate during a time interval of 4.5 ms, what is the magnitude of the induced e.m.f in the loop?

- A 0.026 V      B 0.051 V      C 0.073 V      D 0.098 V

- 24 The uniform wire AB has length 1.0 m and resistance of  $10\ \Omega$ . When NB is 40.0 cm, the a.c. voltmeter reads a steady r.m.s. voltage of 2.5 V.




What is the instantaneous peak power provided by the supply?

- A 0.63 W      B 3.9 W      C 7.8 W      D 16 W
- 25 When monochromatic light of wavelength 440 nm is incident on a metal surface, electrons are emitted. No electrons are emitted from the surface when the wavelength of the incident light is greater than 550 nm.

What is the minimum de Broglie wavelength of an emitted electron?

- A  $3.6 \times 10^{-10}$  m      B  $7.2 \times 10^{-10}$  m      C  $8.1 \times 10^{-10}$  m      D  $1.6 \times 10^{-9}$  m
- 26 The diagram below shows the line spectrum from a hot gas.




Which of the following statements can account for line Y being much brighter than lines X and Z?

- A Line Y has the highest frequency.  
 B Line Y originates in the hottest part of the gas.  
 C Line Y is the result of electrons undergoing transition between two states of greatest energy difference.  
 D Line Y is the result of most electrons undergoing the same transition between the two states involved in the emission.




27 An X-ray spectrum is shown in the diagram below.




What does the value of  $\lambda_{\min}$  represent?

- A The threshold wavelength of the target metal used to produce X-ray.
  - B The wavelength corresponding to the ionization energy of the target metal.
  - C The de Broglie wavelength of the electron with maximum energy.
  - D The wavelength corresponding to all the energy supplied to an electron in the accelerating electric field being converted into a single X-ray photon.
- 28 Two alpha particles with equal energies are fired towards the nucleus of a gold atom.

Which diagram best represents their paths?



- 29 Uranium-235 undergoes fission in a reaction shown below, releasing 195 MeV of energy. The binding energy per nucleon for uranium-235 is 7.6 MeV, and those for caesium and rubidium are approximately **X** MeV.



What is the value of **X**?

- A** 6.7                      **B** 6.8                      **C** 8.4                      **D** 8.5
- 30 Radon  ${}_{86}^{222}\text{Ra}$  is the start of a decay chain that forms bismuth  ${}_{83}^{214}\text{Bi}$  by  $\alpha$  and  $\beta$  emission.

For the decay of each nucleus of radon, how many  $\alpha$ -particles and  $\beta$ -particles are emitted?

	$\alpha$ -particles	$\beta$ -particles
<b>A</b>	1	1
<b>B</b>	2	1
<b>C</b>	1	2
<b>D</b>	2	2

**END OF PAPER**

**Paper 1 Answer Key**

<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>
B	C	C	C	A	B	B	A	B	B
<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>19</b>	<b>20</b>
B	C	A	C	D	D	A	D	B	B
<b>21</b>	<b>22</b>	<b>23</b>	<b>24</b>	<b>25</b>	<b>26</b>	<b>27</b>	<b>28</b>	<b>29</b>	<b>30</b>
A	B	B	D	D	D	D	C	D	B



CANDIDATE  
NAME

--

CIVICS  
GROUP

1	7	-		
---	---	---	--	--

REGISTRATION  
NUMBER

--	--

## PHYSICS

**9749/02**

Paper 2 Structured Questions

**14 September 2018**

**2 hours**

Candidates answer on the Question Paper.  
No Additional Materials are required.

### READ THESE INSTRUCTIONS FIRST

Write your name, civics group and registration number on all the work you hand in.  
Write in dark blue or black pen on both sides of the paper.  
You may use an HB pencil for any diagrams or graphs.  
Do not use paper clips, highlighters, glue or correction fluid.

The use of an approved scientific calculator is expected where appropriate.

Answer **all** questions.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
8	
<b>Total</b>	

This document consists of **22** printed pages and **0** blank pages.

**Data**

speed of light in free space,	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
permeability of free space,	$\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$
permittivity of free space,	$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$ $(1/(36 \pi)) \times 10^{-9} \text{ F m}^{-1}$
elementary charge,	$e = 1.60 \times 10^{-19} \text{ C}$
the Planck constant,	$h = 6.63 \times 10^{-34} \text{ J s}$
unified atomic mass constant,	$u = 1.66 \times 10^{-27} \text{ kg}$
rest mass of electron,	$m_e = 9.11 \times 10^{-31} \text{ kg}$
rest mass of proton,	$m_p = 1.67 \times 10^{-27} \text{ kg}$
molar gas constant,	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
the Avogadro constant,	$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
the Boltzmann constant,	$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$
gravitational constant,	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
acceleration of free fall,	$g = 9.81 \text{ m s}^{-2}$

**Formulae**

uniformly accelerated motion,

$$s = ut + \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

work done on/by a gas,

$$W = p\Delta V$$

hydrostatic pressure,

$$p = \rho gh$$

gravitational potential,

$$\phi = -\frac{Gm}{r}$$

temperature,

$$T/K = T/^\circ\text{C} + 273.15$$

pressure of an ideal gas,

$$p = \frac{1}{3} \frac{Nm}{V} \langle c^2 \rangle$$

mean translational kinetic energy of an ideal gas molecule

$$E = \frac{3}{2} kT$$

displacement of particle in s.h.m.

$$x = x_0 \sin \omega t$$

velocity of particle in s.h.m.

$$v = v_0 \cos \omega t$$

$$= \pm \omega \sqrt{(x_0^2 - x^2)}$$

electric current,

$$I = Anvq$$

resistors in series,

$$R = R_1 + R_2 + \dots$$

resistors in parallel,

$$1/R = 1/R_1 + 1/R_2 + \dots$$

electric potential,

$$V = \frac{Q}{4\pi\epsilon_0 r}$$

alternating current/voltage,

$$x = x_0 \sin \omega t$$

magnetic flux density due to a long straight wire

$$B = \frac{\mu_0 I}{2\pi d}$$

magnetic flux density due to a flat circular coil

$$B = \frac{\mu_0 NI}{2r}$$

magnetic flux density due to a long solenoid

$$B = \mu_0 nI$$

radioactive decay,

$$x = x_0 \exp(-\lambda t)$$

decay constant

$$\lambda = \frac{\ln 2}{t_{\frac{1}{2}}}$$

1 (a) State Newton's three laws of motion.

First Law: .....

.....

.....

Second Law: .....

.....

.....

Third Law: .....

.....

..... [3]

(b) Using the appropriate laws of motion, answer the question for each given situation.

(i) A passenger claimed that he was sitting in the middle of a bus that was moving forward. The driver suddenly applied the brakes and a suitcase that was in the front of the bus flew backwards and hit him.

State and explain if his claim is valid.

.....

.....

.....

..... [2]

(ii) A labourer was tasked with pulling a cart. He reasoned that whatever he exerts on the cart, the cart will exert an equal and opposite force on him. The forces will cancel out and hence it is pointless for him to pull the cart as both he and the cart will not be able to move.

State and explain if his reasoning is correct.

.....

.....

.....

..... [2]

**(iii)** A man is stranded in the middle of a frozen lake with a heavy bag of gold. As there is no friction between him and the surface of the ice, he is unable to crawl back to shore.

State and explain the action that he can take to get back to shore.

.....  
.....  
.....  
.....  
..... [2]

**(c)** A small sedan car makes a head-on collision with a large truck.

**(i)** State and explain if the force experienced by the car is different from that experienced by the truck.

.....  
.....  
..... [1]

**(ii)** Both drivers are securely fastened to their vehicle seats. State and explain if the driver of the truck is likely to experience more severe injuries as compared to the driver of the car.

.....  
.....  
.....  
.....  
..... [2]



2 Fig. 2.1 shows an archer with a bow.




Fig. 2.1

The force  $F$  required to bend the bow and the corresponding displacement  $d$  of the arrow are measured. A plot of  $F$  against  $d$  is shown in Fig. 2.2.




Fig. 2.2

- (a) An experienced archer is able to draw an arrow further back, resulting in a greater displacement of arrow,  $d$ , as compared to a novice archer.

Using Fig. 2.2 and the principle of conservation of energy, suggest an advantage that this extra displacement provides.

.....

.....

.....

.....

..... [2]

- (b) (i) Using Fig. 2.2, estimate the elastic potential energy stored in the bow when an arrow is displaced by 0.50 m.

elastic potential energy = .....J [2]

- (ii) The arrow in (b)(i) is then released by the archer from the bow.

Determine the maximum possible speed of the arrow which has a mass of  $3.5 \times 10^{-2}$  kg.

maximum possible speed = ..... m s<sup>-1</sup> [2]

- 3 (a) Define the *tesla*.

.....

.....

..... [1]

- (b) Charged particles, of speed  $4500 \text{ m s}^{-1}$  and mass  $2.66 \times 10^{-26} \text{ kg}$ , are travelling in a narrow beam in a vacuum as shown in Fig. 3.1.




Fig. 3.1

The charged particles enter a region of uniform magnetic flux density which is 0.200 m wide. The direction of the magnetic flux is pointing out of the paper.

- (i) Using Newton's Law of motion, state and explain the speed of the particles as they exit the uniform magnetic field.

.....

.....

..... [2]

- (ii) Given that the magnitude of the charge on the particles is  $e$ , calculate the radius of the circular motion of the charged particles in the uniform magnetic field.

radius = .....m [2]

- (iii) Hence, calculate the angular velocity of the charged particles in the uniform magnetic field.

angular velocity = ..... rad s<sup>-1</sup> [1]

- (iv) Point **O** is the centre of the circular motion of the charged particles as shown in Fig. 3.1. Show that the angle  $\theta$  is 0.564 rad. [2]

- (v) In another experiment, similar charged particles are now fired into the magnetic field in Fig. 3.1 with different momentum.

- Determine the maximum momentum of particles such that the particles will not exit the magnetic field through XY.

momentum = ..... kg m s<sup>-1</sup> [2]

- Show that the time taken for the particle of mass  $m$  and charge  $q$  in (b)(v)1 to complete a semi-circle within a region of magnetic flux density  $B$  is given by:

$$\frac{\pi m}{Bq} \quad [2]$$

4 (a) State the *principle of superposition*.

.....  
 .....  
 ..... [1]

(b) Noise-cancelling headphones use both Active Noise Control and passive sound-proofing to reduce undesired ambient sounds reaching the ears of the user. In Active Noise Control, a microphone detects ambient noise and a noise-cancellation speaker emits a corresponding “anti-noise” that undergoes destructive interference with the noise.

Suggest the features that the “anti-noise” should have in order to achieve the optimal noise-cancelling effect.

.....  
 .....  
 .....  
 ..... [2]

(c) A student measured the waveforms of the noise and the anti-noise. The following observations were made:

1. There is a 1.0 ms time-lag between the noise and the anti-noise.
2. The amplitudes of the noise and anti-noise were effectively similar.

A sample noise waveform is shown in Fig. 4.1.




Fig. 4.1

(i) On Fig. 4.2, sketch the anti-noise waveform for  $15 \text{ ms} < t < 30 \text{ ms}$ .

[2]




Fig. 4.2

(ii) The final resultant waveform reaching the ears is shown in Fig. 4.3.




Fig. 4.3

Using Fig. 4.3, suggest if Active Noise Control is better suited for noises of higher or lower frequencies.

.....

.....

..... [2]

- 5 (a) You are provided with a galvanometer, a switch, a  $100\ \Omega$  fixed resistor, and some connecting wires.

By completing the potentiometer circuit diagram in Fig. 5.1, explain how the internal resistance of an unknown cell  $\epsilon$  may be determined.




Fig. 5.1

.....

.....

.....

.....

.....

.....

.....

[3]

- (b) Fig 5.2 shows a potentiometer circuit in which the resistance wire is made up of 7 identical strands of unshielded constantan wire.




Fig. 5.2

- (i) Each constantan strand has a diameter of 0.15 mm. Given that XY is 1.00 m long, show that XY has a resistance of 4.0 Ω. (Resistivity of constantan is  $4.90 \times 10^{-7} \Omega \text{ m}$ .) [2]

- (ii) The cells have negligible internal resistances. Explain quantitatively whether a balance length can be found. [3]



- 6 Radiation of wavelength  $4.0 \times 10^{-7} \text{ m}$  is incident on the photo-cathode **C** of a photo-cell. The photo-cell is connected to a potential divider arrangement as shown in Fig. 6.1. The material of **C** has a work function of value  $3.0 \times 10^{-19} \text{ J}$ .




Fig. 6.1

- (a) Explain what is meant by *work function*.

.....  
 .....  
 ..... [1]

- (b) (i) Calculate the maximum kinetic energy of the electrons emitted from the photo-cathode.

maximum kinetic energy = ..... eV [3]

- (ii) Suggest a reason why not all photoelectrons have this maximum energy.

.....  
 .....  
 ..... [1]

- (iii) Calculate the minimum reading of the voltmeter when the milli-ammeter registers no current flowing through it.

voltmeter reading = ..... V [1]

- (c) The sliding contact S is moved from A towards B.

On Fig. 6.2, sketch a graph of current registered by milli-ammeter against potential difference  $V$  measured across the analog voltmeter. [2]




Fig. 6.2

- (d) The polarity of the battery is now reversed. Explain why the milli-ammeter will always register a current flowing through it, regardless of the position of the sliding contact S on AB.

.....  
 .....  
 ..... [2]

- (e) In an actual experiment using the setting in (d), the frequency of the irradiating photons is equal to the threshold frequency. However, there is no photocurrent registered by the milli-ammeter regardless of the position of the sliding contact S on AB.

Suggest a reason for this observation.  
 .....  
 ..... [1]

- 7 (a) For a steady and streamline flow of an incompressible and non-viscous fluid, the total hydraulic head  $H$  of the flow through a particular cross-section of a pipe is given by the expression

$$H = \frac{P}{\rho g} + \frac{v^2}{2g} + h$$

where  $P$  is the pressure of the fluid at the cross-section,  
 $\rho$  is the density of the fluid,  
 $v$  is the velocity of the fluid at the cross-section, and  
 $h$  is the height of the centreline of pipe above a reference level.

$H$  is related the energy in the fluid.

Suggest the type of energy associated with the following terms:

(i)  $\frac{v^2}{2g}$

..... [1]

(ii)  $h$

..... [1]

- (b) Fig. 7.1 shows a steady and fully developed flow through a horizontal pipe of varying diameter. The inlet has an inner pipe diameter  $D_A$  and fluid velocity  $V_A$  while the outlet has an inner pipe diameter  $D_B$  and fluid velocity  $V_B$ .




Fig. 7.1

- (i) Assuming there are no energy losses within the pipe, state how the total hydraulic head at the inlet and at the outlet are related.

..... [1]

- (ii) If the fluid can be considered incompressible, show that  $v_A$  and  $v_B$  are related by the expression

$$\frac{v_A}{v_B} = \frac{D_B^2}{D_A^2} \quad [2]$$

- (iii) The inner diameter of the inlet is 0.100 m and the inner diameter of the outlet is 0.050 m. Water flows into the inlet at a speed of  $8.0 \text{ m s}^{-1}$ .

Using the hydraulic head expression given in (a) and your answers to (b), determine the pressure difference between the inlet and outlet. (Density of water is  $1000 \text{ kg m}^{-3}$ .)

pressure difference = ..... Pa [3]

- (c) In a typical flow between 2 cross-sections of a pipe, there will be energy losses due to frictional forces between the walls of the pipe and the fluid. This is accounted for by including an additional head loss term,

$$H_{\text{Loss}} = f_D \frac{v^2 L}{2gD}$$

where  $f_D$  is the friction factor,  
 $L$  is the length of the pipe, and  
 $D$  is the inner diameter of the pipe.

- (i) State how this additional term will affect your answer to (b)(iii).

.....  
 ....., [1]

- (ii) If the pipe in (b) has a length of 10 m and a friction factor of 0.0038, estimate the magnitude of the difference to your answer in (b)(iii) if frictional losses are considered.

difference = ..... Pa [2]

- (d) A student suggests using water flow in rigid pipes to model blood flow in blood vessels.

- (i) State a difference between water flow in rigid pipes and blood flow in blood vessels.


.....  
.....  
..... [1]

- (ii) The accumulation of fat and cholesterol deposits along the arterial walls has an effect of narrowing the arteries leading to an increase in blood pressure.

State and explain if the modelling proposed by the student will be able to correctly predict this observation.

.....  
.....  
.....  
.....  
.....  
..... [3]

- 8 The International Space Station (ISS) is a habitable artificial satellite. It is maintained at 340 km above the Earth's surface, in what is known as a Low Earth Orbit (LEO). The radius of the Earth is  $6.37 \times 10^6$  m. Fig 8.1 shows how the gravitational field strength of the Earth,  $g$ , varies with distance from the Earth's surface.



**Fig. 8.1**

- (a) Using Fig 8.1, explain why an astronaut experiences apparent weightlessness in the ISS despite a non-zero gravitational field strength at LEO.

.....

.....

.....

..... [2]

- (b) Astronauts in apparent weightlessness may lose muscle mass. Fig. 8.2 shows a model of a spring system which can be used to monitor changes in mass.




Fig. 8.2

It has been suggested that the relationship between the period of oscillation  $T$  and the mass  $M$  is given by:

$$\frac{1}{T^2} = \frac{p}{M} + q$$

where  $p$  and  $q$  are constants.

Fig. 8.3 shows the experimental results obtained using a stopwatch to measure the time taken for oscillations.

$M / \text{kg}$	time taken for 20 oscillations		$T / \text{s}$	$\frac{1}{T^2} / \text{s}^{-2}$	$\frac{1}{M} / \text{kg}^{-1}$
	$t_1 / \text{s}$	$t_2 / \text{s}$			
0.100	10.34	10.38	0.5180	3.727	10.0
0.150	12.07	11.93	0.6000	2.778	6.67
0.200	13.39	13.43	0.6705	2.242	5.00
0.250	14.34	14.32	0.7165	1.948	4.00
0.300	15.23	15.23	0.7615	1.724	3.33
0.350	15.70	15.64	0.7835	1.629	2.86

Fig. 8.3

The corresponding values of  $\frac{1}{T^2}$  and  $\frac{1}{M}$  are plotted on the graph in Fig. 8.4.




Fig. 8.4

(i) Using Fig. 8.4, determine  $p$  and  $q$ .

$p = \dots\dots\dots$

$q = \dots\dots\dots$  [3]



- (ii) It has been suggested that the variation of period,  $\partial T$ , is related to an astronaut's mass loss by  $\partial T = \frac{2pT^3}{M} \times (\text{fractional mass loss})$ .

Determine  $\partial T$  corresponding to an initial mass of  $M = 0.5$  kg and a fractional mass loss of 0.10.

$$\partial T = \dots\dots\dots\text{s [2]}$$

- (iii) According to medical opinion, a muscular mass loss of 10% is concerning. Taking reference from the data available, discuss if the experiment is able to detect the variation  $\partial T$  for a mass as small as 0.5 kg.

.....  
 .....  
 ..... [2]

**END OF PAPER**



## PHYSICS

9749/02

Paper 2 Structured Questions  
MARK SCHEME  
Maximum Mark: 80


14 September 2018

Question	Marks
1(a)	
N1L: An object at rest will remain <u>at rest</u> and an object in motion will remain in motion at <u>constant velocity</u> in the <u>absence of an external resultant force</u> .	B1
N2L: The <u>rate</u> of change of momentum of a body is <u>directly</u> proportional to the <u>resultant</u> force acting on the body and occurs in the direction of the resultant force.	B1
N3L: If body A exerts a force on body B, then body B exerts a force of the <u>same type</u> that is equal in magnitude and opposite in direction on body A.	B1
1(b)(i)	
Initially, suitcase moving with <b>same velocity as bus</b> before braking	B1
By N1L, suitcase will continue to move forward in the absence of external force acting on it because not secured (or small frictional force between when the suitcase was still in contact with the bus that is not sufficient to stop it from moving forward) when brakes applied	B1
Claim invalid	A0
1(b)(ii)	
The two equal and opposite forces act on <u>cart and labourer separately</u>	B1
Cart pulled forward by the force applied by the labourer on the cart.	B1
Labourer moved forward by the <u>frictional force between feet/shoes and ground</u> .	B1
Reasoning false	A0
1(b)(iii)	
Throw bag of gold in direction away from shore	B1
By N3L, man experiences a force equal in magnitude and opposite in direction from the force he exerted on the bag of gold. This forces accelerates the man towards shore during pushing, thereafter he will glide over ice at constant speed	B1
1(c)(i)	
By N3L, force experienced by car <u>same</u> as force experienced by truck	B1
1(c)(ii)	
Total mass of truck and its driver <u>larger than</u> total mass of car and its driver, hence the truck and its driver experiences smaller change in velocity over same duration of time	B1
Safety belt exert smaller force on truck driver in stopping his forward velocity	B1
Truck driver likely less injured than car driver	A0

Question	Marks
<b>2(a)</b> Area under F-x graph is w.d. in stretching bow. More potential energy is stored	B1
Arrows gain more KE, has less deviation from intended flight path <b>OR</b> Arrows gain more KE, has farther range	B1
<b>2(b)(i)</b> By counting squares under F-x graph, Elastic potential energy = area under F-x graph = $1.25 \text{ J} \times 64.5$ ( $\pm 2$ squares) = 81 J (78.1 J to 83.1J)	M1
<b>2(b)(ii)</b> By Conservation of Energy, Loss in elastic PE = Gain in KE for arrow $81 = \frac{1}{2}(3.5 \times 10^{-2})v^2$ $v = 68.0 \text{ m s}^{-1}$ (66.8 J to 68.9 J)	M1
<b>3(a)</b> The tesla is the uniform magnetic flux density which when acting at right angle to a straight conductor carrying a current of 1 ampere produces a force per unit length of 1 Newton per metre on the conductor.  Or  If a conductor carrying a current of 1 ampere is placed at right angles to a uniform magnetic field of flux density 1 Tesla, then the force per unit length on the conductor is 1 newton per metre.	B1
<b>3(b)(i)</b> Charged particles moving perpendicular to a magnetic field will experience a resultant magnetic force perpendicular to its motion. Hence no work is done. By Newton's 2 <sup>nd</sup> Law, the acceleration of the particles is in the same direction as the resultant force. The direction of the particles changes but not its speed. By Newton's 1 <sup>st</sup> law, upon exit, the particles will move in a straight line with a speed of $4500 \text{ ms}^{-1}$ .	B1
<b>3(b)(ii)</b> Magnetic force provides centripetal force for particle's circular motion $F = Bqv = \frac{mv^2}{r}$ $r = \frac{mv}{Bq}$ $= \frac{(2.66 \times 10^{-26})(4500)}{(2 \times 10^{-3})(1.6 \times 10^{-19})}$ $= 0.374 \text{ m}$	B1
<b>3(b)(iii)</b> $v = r\omega$ $\omega = \frac{v}{r} = \frac{4500}{0.374}$ $= 12030 \text{ rad s}^{-1}$	M1
	A1

Question	Marks
<b>3(b)(iv)</b> $\sin \theta = \frac{OX}{r}$ $\theta = \sin^{-1} \left( \frac{0.2}{0.374} \right)$ $= 0.564 \text{ rad}$	M1  A1
<b>3(b)(v)1.</b> $r = \frac{mv}{Bq} = \frac{P}{Bq}$ $P = rBq = (0.2)(2 \times 10^{-3})(1.6 \times 10^{-19})$ $= 6.4 \times 10^{-23} \text{ kg m s}^{-1}$	M1  A1
<b>3(b)(v)2.</b> magnetic force F provides centripetal force for the particle's circular motion $F = Bqv = mr\omega^2$ $Bq(r\omega) = mr\omega^2$ $Bq = m \left( \frac{2\pi}{T} \right)$ $T = 2\pi \frac{m}{Bq}$ For semicircle, $t = \frac{T}{2} = \frac{\pi m}{Bq}$	M1        A1


- 4(a)** Two or more waves of the same kind meet at a point at the same time, displacement of the resultant wave is vector sum of the displacements of the individual waves at that point at that time. B1
- 4(b)** constant phase difference of  $\pi$  (rad)  
same amplitude B1  
B1
- 4(c)(i)** Amplitude



- phase difference (shown by reflection in x-axis) B1  
time lag (shown by translation of 2 small squares to right) B1

Question		Marks
4(c)(ii)	<u>longer durations of destructive interference</u> for noise waves of <u>longer periods</u> ; correspond to <u>lower frequencies</u>	M1 A1

5(a)



- Correct drawing with switch in the correct position. B1

Turn off the switch. Find balance length when galvanometer shows null reading. B1

Find  $\epsilon$  using  $\epsilon = V_{AC} = kL_{AC}$  and  $V_{AB} = E = kL_{AB}$

Turn on the switch.

Find the new balance length,  $L_{AC}'$

B1

To find  $r$ :

Compare  $V_{AC}' = kL_{AC}'$  and  $V_{AB} = E = kL_{AB}$

$V_{AC}' = kL_{AC}' = (R/R+r) \cdot \epsilon$

Solve  $r$ .

Question	Marks
<p>5(b)(i) For single strand:</p> $R = \frac{\rho L}{A}$ $= \frac{(4.9 \times 10^{-7})(1)}{\pi \left( \frac{0.15 \times 10^{-3}}{2} \right)^2}$ <p>(= 27.7 <math>\Omega</math>)</p>	M1
<p>For 7 strands:</p> $\frac{1}{R_{eff}} = 7 \left( \frac{1}{R} \right)$ $R_{eff} = \frac{R}{7}$ $= \frac{(4.9 \times 10^{-7})(1)}{\pi(7) \left( \frac{0.15 \times 10^{-3}}{2} \right)^2}$ <p>= 3.96 <math>\Omega</math></p> <p>= 4.0 <math>\Omega</math></p>	A1
<p>5(b)(ii) Potential difference across the 7 <math>\Omega</math> resistor = <math>\frac{7}{14+7} \times 12 = 4 V</math></p>	B1
<p>By potential divider rule, potential difference across 0.9 m potentiometer</p> $= \frac{0.9 \times 3.96}{3.96+10} \times 12 = 3.06 V$	B1
<p>Since the potential difference across the potentiometer is less than the potential difference across the 7 <math>\Omega</math> resistor, balance length cannot be achieved.</p>	B1
<p>6(a) Work function is the minimum amount of energy required for an electron to escape from the surface of a metal.</p>	B1
<p>6(b)(i) Using conservation of energy,</p>	
$KE_{max} = hf - \phi$ $= \frac{6.63 \times 10^{-34} \times 3.0 \times 10^8}{4.0 \times 10^{-7}} - 3.0 \times 10^{-19}$ $= 1.97 \times 10^{-19} \text{ J}$ $= \frac{1.97 \times 10^{-19}}{1.60 \times 10^{-19}} = 1.23 \text{ eV}$	M1
<p>6(b)(ii) Some of the electrons in material C may <u>need energy greater than the work function energy to be liberated as photoelectrons as they are beneath the metal surface</u>. Hence, not all photoelectrons have this maximum kinetic energy.</p>	B1

Question	Marks
6(b)(iii) p.d. = 1.23 V	B1
6(c)	
Correct shape of curve (Only in 1 quadrant of graph) Stopping potential clearly and correctly labelled numerically.	B1 B1
6(d) When the polarity is being reversed, <u>C will always be at a lower potential compared to D</u> . Photoelectrons released from C will therefore <u>always be attracted to D</u> and close the circuit for current to flow through the milliammeter.	M1 A1
6(e) Metal surface is oxidized. <b>OR</b> The intensity of the radiation is not sufficiently high enough to liberate enough photoelectrons for them to be measureable by the milliammeter.	B1
Any one of the above points.	

---

7(a)(i)	kinetic energy (of the fluid)	B1
7(a)(ii)	gravitational potential energy (of the fluid)	B1
7(b)(i)	They are the same/ equal.	B1
7(b)(ii)	mass flow (rate) at inlet = mass flow (rate) at outlet	B1

$$\frac{m_A}{t} = \frac{m_B}{t}$$

$$\frac{\rho \frac{\pi D_A^2}{4} l_A}{t} = \frac{\rho \frac{\pi D_B^2}{4} l_B}{t}$$

$$D_A^2 v_A = D_B^2 v_B$$

$$\frac{v_A}{v_B} = \frac{D_B^2}{D_A^2}$$

B1

Question	Marks	
7(b)(iii)		
$\frac{v_{out}}{v_{in}} = \frac{D_{in}^2}{D_{out}^2}$		
$v_{out} = \left(\frac{0.100}{0.050}\right)^2 v_{in}$ $= 4v_{in}$	C1	
$\frac{P_{in}}{\rho g} + \frac{v_{in}^2}{2g} + h_{in} = \frac{P_{out}}{\rho g} + \frac{v_{out}^2}{2g} + h_{out}$	M1	
$\frac{P_{out} - P_{in}}{\rho g} = \frac{v_{in}^2 - v_{out}^2}{2g}$		
$\Delta P = \frac{\rho(v_{in}^2 - v_{out}^2)}{2}$		
$= \frac{\rho(v_{in}^2 - (4v_{in})^2)}{2}$		
$= -480 \text{ kPa}$	A1	
7(c)(i)	The pressure difference will be higher.	B1
7(c)(ii)	$v_{ave} = 2.5v_{in}$ $D_{ave} = 0.075 \text{ m}$ (or any other appropriate averaging done)	C1
	$p_{Loss} = H_{Loss} \times \rho g$ $= f_D \frac{v_{ave}^2 L}{2gD_{ave}} \times \rho g$ $= 0.0038 \times \frac{(2.5 \times 8)^2 \times 10}{2 \times 0.075} \times 1000$ $= 101 \text{ kPa}$	A1
7(d)(i)	<ul style="list-style-type: none"> <li>✓ Blood vessels are flexible</li> <li>✓ (Higher) friction factor / viscosity of blood</li> <li>✓ Blood is a suspension rather than a pure fluid.</li> <li>✓ Capillary action due to the small diameter of blood vessels.</li> </ul>	B1
7(d)(ii)	As the arteries narrow, the model predicts that the <u>velocity of the flow</u> at this cross section will be <u>higher</u> . From the expression in (a), the <u>pressure of the blood</u> at the narrow cross-section will be <u>lower</u> . Thus, this model proposed by the student <u>will not be able to correctly predict</u> this observation.	M1
		M1
		A1



Question	Marks
<b>8(a)</b> gravitational force provides (just enough) centripetal acceleration of $8.825 \text{ m s}^{-2}$ on both ISS and astronaut to keep them in circular orbit around Earth	
OR	
both ISS and astronaut experience $8.825 \text{ ms}^{-2}$ of acceleration directed to the centre of the earth due to gravity	M1
no contact force by ISS on astronaut.	A1
<b>8(b)(i)</b>	
best fit straight line with line thickness not comparable to half sq $p = 0.298 \text{ kg s}^{-2}$ $q = 0.759 \text{ s}^{-2}$	B1 B1 B1
<b>8(b)(ii)</b>	
For $\frac{1}{T^2} = \frac{p}{M} + q \rightarrow T = \sqrt{\left(\frac{p}{M} + q\right)^{-1}}$	
For $M = 0.5 \text{ kg}$ , $T = \sqrt{\left(\frac{0.298}{0.5} + 0.759\right)^{-1}} = 0.859 \text{ s}$	C1
$\partial T = \frac{2(0.298)(0.858)^3}{0.5} \times (10\%) = 0.08 \text{ s (1 s.f.)}$	A1
<b>8(b)(iii)</b>	
Yes. Period has more precision that what is required of variation	B1
from table of values (Fig. 8.3), the period is to 4 decimal points while the variation which is to 2 d.p.	B1



CANDIDATE  
 NAME

--

CIVICS  
 GROUP

1	7	-	
---	---	---	--

REGISTRATION  
 NUMBER

--	--

## PHYSICS

**9749/03**

Paper 3 Longer Structured Questions

**18 September 2018**

**2 hours**

Candidates answer on the Question Paper.  
 No Additional Materials are required.

### READ THESE INSTRUCTIONS FIRST

Write your name, civics group and registration number on all the work you hand in.  
 Write in dark blue or black pen on both sides of the paper.  
 You may use an HB pencil for any diagrams or graphs.  
 Do not use paper clips, highlighters, glue or correction fluid.

The use of an approved scientific calculator is expected where appropriate.

#### Section A

Answer **all** questions.

#### Section B

Answer **one** question only

You are advised to spend one and a half hours on Section A and half an hour on Section B

At the end of the examination, fasten all your work securely together.  
 The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
<b>1</b>	
<b>2</b>	
<b>3</b>	
<b>4</b>	
<b>5</b>	
<b>6</b>	
<b>7</b>	
<b>8</b>	
<b>Total</b>	

This document consists of **25** printed pages and **1** blank page.

**Data**

speed of light in free space,	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
permeability of free space,	$\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$
permittivity of free space,	$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$ $(1/(36 \pi)) \times 10^{-9} \text{ F m}^{-1}$
elementary charge,	$e = 1.60 \times 10^{-19} \text{ C}$
the Planck constant,	$h = 6.63 \times 10^{-34} \text{ J s}$
unified atomic mass constant,	$u = 1.66 \times 10^{-27} \text{ kg}$
rest mass of electron,	$m_e = 9.11 \times 10^{-31} \text{ kg}$
rest mass of proton,	$m_p = 1.67 \times 10^{-27} \text{ kg}$
molar gas constant,	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
the Avogadro constant,	$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
the Boltzmann constant,	$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$
gravitational constant,	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
acceleration of free fall,	$g = 9.81 \text{ m s}^{-2}$

**Formulae**

uniformly accelerated motion,

$$s = ut + \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

work done on/by a gas,

$$W = p\Delta V$$

hydrostatic pressure,

$$p = \rho gh$$

gravitational potential,

$$\phi = -\frac{Gm}{r}$$

temperature,

$$T/K = T/^\circ\text{C} + 273.15$$

pressure of an ideal gas,

$$p = \frac{1}{3} \frac{Nm}{V} \langle c^2 \rangle$$

mean translational kinetic energy of an ideal gas molecule

$$E = \frac{3}{2}kT$$

displacement of particle in s.h.m.

$$x = x_0 \sin \omega t$$

velocity of particle in s.h.m.

$$v = v_0 \cos \omega t$$

$$= \pm \omega \sqrt{(x_0^2 - x^2)}$$

electric current,

$$I = Anvq$$

resistors in series,

$$R = R_1 + R_2 + \dots$$

resistors in parallel,

$$1/R = 1/R_1 + 1/R_2 + \dots$$

electric potential,

$$V = \frac{Q}{4\pi\epsilon_0 r}$$

alternating current/voltage,

$$x = x_0 \sin \omega t$$

magnetic flux density due to a long straight wire

$$B = \frac{\mu_0 I}{2\pi d}$$

magnetic flux density due to a flat circular coil

$$B = \frac{\mu_0 NI}{2r}$$

magnetic flux density due to a long solenoid

$$B = \mu_0 nI$$

radioactive decay,

$$x = x_0 \exp(-\lambda t)$$


decay constant

$$\lambda = \frac{\ln 2}{t_{\frac{1}{2}}}$$

**Section A**


Answer **all** the questions in this Section in the spaces provided.

- 1 A spring is attached at one end to a fixed point and hangs vertically with a cube attached to the other end. The cube is initially supported at a height  $h$  above the water surface such that the spring is at its natural length, as shown in Fig. 1.1.



**Fig. 1.1**

The cube has a weight of 4.0 N and sides of length 5.1 cm. The cube is lowered gently into water. The cube reaches equilibrium with its base at a depth of 7.0 cm below the water surface, as shown in Fig. 1.2. The density of the water is  $1000 \text{ kg m}^{-3}$ .



**Fig. 1.2**

- (a) Calculate the difference in the pressure exerted by the water on the bottom face and on the top face of the cube.

difference in pressure = ..... Pa [2]

(b) Using your answer in (a), show that the upthrust acting on the cube is 1.3 N. [1]

(c) Determine the force exerted on the spring by the cube when it is in equilibrium in the water.

force = ..... N [2]

(d) The spring obeys Hooke's law and has a spring constant of  $30 \text{ N m}^{-1}$ .

Determine the height  $h$ .

$h = \dots\dots\dots \text{ m}$  [3]

(e) The cube in the water is suddenly detached from the spring.

(i) Determine the initial acceleration of the cube.

initial acceleration = .....  $\text{m s}^{-2}$  [2]

(ii) Describe and explain the variation of the acceleration of the cube as it sinks in the water.

.....

.....

..... [2]

2 (a) State what is meant by *simple harmonic motion*.

.....

.....

..... [2]

(b) A small frictionless trolley is attached to a fixed point **P** by means of a spring. A second spring is used to attach the trolley to a variable frequency oscillator, as shown in Fig. 2.1.




Fig. 2.1

Both springs remain extended within their limits of proportionality. Initially, the oscillator is switched off. The trolley is displaced horizontally along the line joining the two springs and is then released. The variation with time  $t$  of the velocity  $v$  of the trolley is shown in Fig. 2.2.




Fig. 2.2

(i) Using Fig. 2.2, state two different times at which

1. the displacement of the trolley is zero,

time = ..... s and ..... s [1]

2. the acceleration in one direction is maximum.

time = ..... s and ..... s [1]

(ii) Determine the frequency of oscillation of the trolley.

frequency = ..... Hz [1]

(iii) The variation with time of the displacement of the trolley is sinusoidal. The variation with time of the velocity of the trolley is also sinusoidal.

State the phase difference between the displacement and the velocity.

phase difference = ..... ° [1]

(c) The oscillator is now switched on. The amplitude of variation of the oscillator is constant. The frequency  $f$  of vibration of the oscillator is varied. The trolley is forced to oscillate by means of vibrations of the oscillator.

(i) Distinguish between *free oscillations* and *forced oscillations*

.....  
.....  
.....  
.....  
..... [2]



- (ii) It was observed that the trolley vibrates with different amplitudes as the frequencies of the oscillator changes.

On the axis provided on Fig. 2.3, sketch a possible amplitude-frequency graph for this trolley. [1]



Fig. 2.3

- (iii) State the approximate frequency at which the amplitude is maximum.

frequency = ..... Hz [1]

- (iv) The amplitude of the oscillations may be reduced without changing significantly the frequency at which the amplitude is a maximum.

State and explain how this may be done. You may draw on Fig. 2.1 to support your answer.

.....

.....

.....

..... [2]

3 (a) Define *electric potential* at a point.

.....  
 .....  
 ..... [1]

(b) Two similar positive point charges of magnitude  $e$  are set up along the line  $XY$  as shown in Fig. 3.1.




Fig. 3.1

(i) On Fig 3.2, sketch how the electric potential varies over  $XY$ . [2]




Fig. 3.2

(ii) Describe how the electric force acting on a known charge  $q$  placed at a point along  $XY$  could be obtained from Fig 3.2.

.....  
 ..... [2]

(c) (i) A negative point charge of  $-e$  is now placed at the midpoint of  $XY$ .

State the magnitude of electric force experienced by the point charge

electric force = ..... N [1]

- (ii) The negative point charge at the midpoint in XY in (c)(i) is moved to the position Z as shown in Fig 3.3.




Fig. 3.3

Within the bounded region of Fig 3.4, sketch the resultant electric field lines of the 3 point charges. [3]




Fig. 3.4

- (iii) Determine the work done by the electric field to completely separate the 3 point charges in Fig 3.3.

work done = ..... J [3]

4 (a) State the *First Law of Thermodynamics*.

.....

.....

..... [1]

(b) 0.050 moles of an ideal gas is contained within an uninsulated cylinder with a movable piston as shown in Fig. 4.1.




Fig. 4.1

The piston moves slowly outwards, resulting in the variation of pressure shown in Fig. 4.2




Fig. 4.2

(i) The process A to B occurs along an isotherm.

Determine the temperature along this isotherm.

temperature = ..... K [2]

- (ii) Calculate the total kinetic energy as a result of the random motion of the gas molecules in the cylinder.

total kinetic energy = ..... J [1]

- (iii) Estimate the amount of work done by gas as it expands from A to B

work done = ..... J [2]

- (iv) A student states that no heat flows into or out of the gas during the process A to B as the temperature of the gas did not change.

State and explain the validity of his statement.

.....  
 .....  
 .....  
 ..... [3]

- (c) In another set up with the same starting point as the one shown in Fig. 4.1, the piston is moved very quickly outwards expanding the gas to a volume of  $1.2 \times 10^{-3} \text{ m}^3$ . As a result, the temperature of the gas decreases to 144 K.

On Fig. 4.2, sketch the variation with volume of the pressure of the gas. [2]

- 5 Fig. 5.1 shows the structure of a geophone which is used by geophysicists to determine the speed of seismic waves traveling within the ground layer.




Fig. 5.1

The spike of the geophone is inserted into the ground. When a seismic vibration moves the case and coil, the magnet remains stationary due to its inertia.

- (a) State Faraday's law of electromagnetic induction.

.....  
 ..... [1]

- (b) Using Faraday's law, explain how an e.m.f. is generated between the terminals when seismic waves pass through the ground.

.....  
 .....  
 .....  
 ..... [2]

(c) A coil of 50 turns generates a maximum e.m.f. of 85 mV in a geophone.

Calculate the rate of change of magnetic flux needed to generate this e.m.f.

rate of change of magnetic flux = .....  $\text{Wb s}^{-1}$  [2]

(d) Suggest 2 changes to the geophone which will make it more sensitive to the seismic vibrations.

1. .... [1]

2. .... [1]

6 In the hairdryer shown in Fig. 6.1, an alternating current passes through the heating coil when the switch is closed.




Fig. 6.1

(a) By reference to heating effect, explain what is meant by the root-mean-square (r.m.s) value of an alternating current.

..... [1]



**(b)** The hair dryer is connected to the mains supply of 120 V r.m.s. and frequency 50 Hz. The heating coil delivers a power of 1200 W.

**(i)** Calculate the peak current through the heating coil.

peak current = ..... A [2]

**(ii)** Write an equation, in terms of the elapsed time  $t$ , for the current that passes through the heating coil, given that the instantaneous power output is zero at  $t = 0$  s. [1]

**(c)** The primary coil of a transformer is connected to a 2.4 kV r.m.s. supply. The secondary coil is connected to the hair dryer and the current flowing through the heating coil has the same value as that calculated in **(b)(i)**.

The transformer is non-ideal and electrical energy is converted to thermal energy in the transformer at a rate of 600 W.

Determine the r.m.s. current in the primary coil.

r.m.s. current = ..... A [2]

**Section B**

Answer any **one** question in this Section in the spaces provided.


7 (a) (i) Explain what is meant by an *alternating current*.

.....

.....

..... [1]

(ii) Fig. 7.1. shows a metal wire held taut between a knife edge X and a smooth pulley P. The wire passes between opposing poles of permanent bar magnets.



**Fig. 7.1**

The wire vibrates when a sinusoidal alternating source is connected across the wire. Explain how these vibrations are created. Describe the properties of these vibrations.

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [4]

**(iii)** When the frequency of the alternating source is 120 Hz, a standing wave wire in its fundamental mode is observed in the wire. XP is 2.0 m in length.

1. Calculate the speed of the wave in the wire.

speed of wave = ..... m s<sup>-1</sup> [2]

2. Explain, with reference to the formation of a stationary wave, what is meant by the speed calculated in **(a)(iii)1**.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

[4]

- (iv) When the mass hanging below pulley P is doubled, the fundamental mode standing wave is observed to occur when the frequency of the alternating source is 170 Hz. When the mass was triple that of the original set up, the fundamental mode frequency is 208 Hz.

Suggest if the relationship

$$v \propto \sqrt{T}$$


is valid, where  $v$  is the wave velocity in the metal wire, and  
 $T$  is the tension in the metal wire.

[3]

- (b) (i) State what is meant by the *diffraction* of a wave.

.....  
..... [1]

- (ii) Light of wavelength 633 nm from a laser is directed normally at a diffraction grating, as illustrated in Fig. 7.2.



**Fig. 7.2**

The diffraction grating is situated at the centre of a circular scale, marked in degrees. The readings on the scale for the second order diffracted beams are 160° and 188°.

Calculate the number of lines per unit length of the slits in the diffraction grating.

number of lines per unit length = ..... m<sup>-1</sup> [4]

- (iii) Suggest why the non-central fringes produced by light passing through a diffraction grating is brighter than that from the same source with a double slit.

.....  
 ..... [1]

- 8 (a) In an  $\alpha$ -particle scattering experiment, an  $\alpha$ -particle is travelling in a vacuum towards the centre of a gold nucleus, as illustrated in Fig. 8.1.




Fig. 8.1

The gold nucleus has a charge  $79e$ . At a large distance from the gold nucleus, the  $\alpha$ -particle has energy  $7.7 \times 10^{-13} \text{ J}$ .

- (i) The  $\alpha$ -particle does not collide with the gold nucleus.

Show that the radius of the gold nucleus must be less than  $4.7 \times 10^{-14} \text{ m}$ . [2]

- (ii) The results of the  $\alpha$ -particle scattering experiment provide evidence for the structure of the atom.

result 1: The vast majority of  $\alpha$ -particles pass straight through the metal foil or are deviated by small angles.

result 2: A very small minority of  $\alpha$ -particles are scattered through angles greater than  $90^\circ$  and up to  $180^\circ$ .

State what may be inferred from

1. result 1,

.....  
 ....., [1]

2. result 2.


.....

.....

.....

..... [2]

(b) One nuclear reaction that can take place in a nuclear reactor may be represented, in part, by the equation



Data for a nucleus and some particles are given in Fig. 8.2

nucleus or particle	mass / u
${}_{57}^{139}\text{La}$	138.955
${}_0^1\text{n}$	1.00863
${}_1^1\text{p}$	1.00728
${}_{-1}^0\text{e}$	0.000549

Fig. 8.2

(i) Complete the nuclear reaction shown above. [1]

(ii) Calculate the binding energy per nucleon, in MeV, of lanthanum-139 ( ${}_{57}^{139}\text{La}$ ).

binding energy per nucleon = ..... MeV [3]

(iii) State and explain whether the binding energy per nucleon of uranium-235 ( $^{235}_{92}\text{U}$ ) will be greater, equal to or less than your answer to (b)(ii).

.....

.....

.....

..... [2]

(c) A radiation detector is placed close to a radioactive source as shown in Fig. 8.3.




Fig. 8.3

The emissions from the radioactive source include both  $\beta$ -particles and  $\gamma$ -ray photons. The  $\beta$ -particles emitted have energies up to a maximum of 0.61 MeV.

(i) Suggest two reasons why the activity of the source and the measured count rate may be different.

1. .... [1]

2. .... [1]



(ii) Explain why the emitted  $\beta$ -particles have a range of energies.

.....  
 ..... [1]

(iii) The  $\gamma$ -ray photons emitted have specific energies. Suggest why this is so.

.....  
 ..... [1]

(d) The variation with time  $t$  of the measured count rate in (c) is as shown in Fig. 8.4.




Fig. 8.4

(i) Use Fig. 8.4 to determine the half-life of the radioactive source.

half-life = ..... hours [3]

(ii) The readings in Fig. 8.4 were obtained at room temperature.

A second sample of this radioactive source is heated to a temperature of 500 °C. The initial count rate at time  $t = 0$  is the same as that in Fig. 8.4. The variation with time  $t$  of the measured count rate from the heated source is determined.

State and explain if there are any differences in

1. the half-life,

.....  
..... [1]

2. the measured count rate for any specific time.

.....  
..... [1]

**END OF PAPER**

**BLANK PAGE**



## PHYSICS

9749/03

Paper 3 Longer Structured Questions

18 September 2018

MARK SCHEME

Maximum Mark: 80


Question		Marks
1(a)	Difference in pressure $\Delta p = \rho g \Delta h$ $= 1000 \times 9.81 \times 0.051$ $= 500.31 \text{ Pa}$ $= 500 \text{ Pa}$	M1 A1
1(b)	Upthrust $U = \Delta p \times 5.1 \times 5.1 \times 10^{-4}$ $= 1.3005 = 1.3 \text{ N}$	M1 A0
1(c)	$U + T = W$ $1.3 + T = 4.0$ $T = 2.7 \text{ N}$ By N3L Force on spring by cube is 2.7 N downwards.	M1 A1
1(d)	Let the extension of the spring be $x$ . $T = k x$ $2.7 = 30 x$ $x = 0.090 \text{ m} = 9.0 \text{ cm}$ initial height above surface, $h$ $= 9.0 - 7.0$ $= 2.0 \text{ cm}$ $= 0.020 \text{ m}$	M1 A0 M1 A1
1(e)(i)	$2.7 = 4.0/9.81 \times a$ $a = 6.62 \text{ m s}^{-2}$ (Downwards)	M1 A1
1(e)(ii)	Viscous force is proportional to speed. Resultant force downwards = Weight – Viscous Force (- Upthrust if considered not negligible) $a$ decreases as velocity increases. When viscous force = weight, $a = 0 \text{ m s}^{-2}$	B1 B1

Question		Marks
2(a)	acceleration / force proportional to displacement (from a fixed point)	B1
	either acceleration and displacement in opposite directions or acceleration always directed towards a fixed point	B1
2(b)(i)1.	0.1 s, 0.3 s, 0.5 s, etc (any two)	B1
2(b)(i)2.	either <b>0 s, 0.4 s, 0.8 s, 1.2 s</b>  or <b>0.2 s, 0.6 s, 1.0 s</b> (any two)	B1
2(b)(ii)	period = 0.4 s frequency = 2.5 Hz	B1
2(b)(iii)	phase difference = 90°	B1
2(c)	free: (body oscillates) without any loss of energy / no resistive forces / no external forces applied	B1
	forced: continuous energy input (required) / body is made to vibrate by an (external) periodic force / driving oscillator	B1
2(d)(i)	<p>Amplitude of forced oscillations</p> <p>natural frequency of the oscillator</p>	
	Any of these shapes and max peak at natural frequency	B1
2(d)(ii)	frequency = 2.4 - 2.5 Hz	B1
2(d)(iii)	e.g. (1) attach sheet of card to trolley bigger than the cross section of the trolley	M1
	increases damping / frictional force	A1
	e.g. (2) reduce oscillator amplitude	M1
	reduce power / energy input to system	A1

**Question****Marks**

**3(a)** the work done per unit positive charge by an external agent in bringing a small test charge from infinity to that point, without any change in the kinetic energy of the charge.

**3(b)(i)**




B1  
B1

**3(b)(ii)** The electric field strength ( $E$ ) at a point between X and Y can be found by finding the negative of the gradient of the tangent drawn at that point on the graph in 3(b)(i). B1

The electric force acting on the known charge  $q$  can be found by multiplying the electric strength  $E$  at that point with the known charge  $q$ . B1

**3(c)(i)** Zero



Electric field lines from positive charge to negative charge shown with arrows. B1

- Lines do not intersect B1
- Line start and end with charge B1
- Line are closer near the centre line B1
- Symmetrical in shape

Question	Marks
----------	-------

**3(c)(iii)**     To move the positive charge q at X to infinity:  
 Total electric potential at X due to the charges at Z and Y,  $V_x = 0$      B1  
 Work done by external agent to move +e from X to infinity,  $W_x = e.(0-V_x)$   
 $= e.0$   
 $= 0$

To move - e at Z to infinity:  
 Total electric potential at Z ( $V_z$ ) is due to the electric charge at Y =  $+e/4\pi\epsilon_0(0.06)$      B1

Work done by external agent to move -e from Z to infinity,  $W_z = -e.(0-V_z)$   
 $= -e. (0 - e/4\pi\epsilon_0(0.06))$   
 $= e^2/4\pi\epsilon_0(0.06)$   
 $= 3.837 \times 10^{-27} \text{ J}$

No work is needed to move the last charge at Y to infinity.

Hence total work done by external agent to separate the 3 charges completely =  $0+3.837 \times 10^{-27} +0$      B1  
 $= 3.84 \times 10^{-27} \text{ J}$

Work done by electric field to separate the 3 charges totally from one another  
 $= - 3.84 \times 10^{-27} \text{ J}$

**OR**

Other variation of method

Work done by external agent to assemble charges one at a time  
 $= -$  work done by external agent to separate the 3 charges totally from one another  
 $=$  work done by electric field to separate the 3 charges totally from one another.

**4(a)**     The increase in the internal energy of a system is the sum of the external work done on the system and the heat supplied to the system.     B1

**4(b)(i)**     Using  $T = \frac{pV}{nR}$      M1

Taking any point:

Point A:  $T_A = \frac{(400 \times 10^3)(0.3 \times 10^{-3})}{(0.050)(8.31)} = 289 \text{ K}$

Point B:  $T_B = \frac{(100 \times 10^3)(1.2 \times 10^{-3})}{(0.050)(8.31)} = 289 \text{ K}$

Therefore, temperature is 289 K.     A1

**4(b)(ii)**     Kinetic energy =  $\frac{3}{2}nRT = \left(\frac{3}{2}\right)(0.050)(8.31)(289) = 180 \text{ J}$      M1  
 -     A0



Question	Marks
----------	-------


**4(b) (iii)** Work done by gas = area under the  $p$ - $V$  graph  
 = (25 squares+9 missing squares)  $(50 \times 10^3) (0.10 \times 10^{-3})$   
 = 170 J (allow 165 to 175J) M1  
A1

**4(b) (iv)** temperature unchanged, no increase in the internal energy.  $\Delta U = 0J$   
 When the gas expands from A to B, work done on the gas is  $w = - 170 J$ . M1

Using 1<sup>st</sup> Law of Thermodynamics:  $\Delta U = w + q$   
 $0 = - 170 + q$   
 Heat supplied,  $q = +170 J$  A1

$q$  is positive  $\rightarrow$  heat is supplied to the gas during the expansion of the gas to ensure that there is no increase in the internal energy of the gas. Therefore, the student claim is false.

**4(c)**



Downwards sloping curve below AB B1  
 Ends at (1.2, 50) B1

**5(a)** The induced e.m.f. is directly proportional to the rate of change of magnetic flux linkage. B1

**5(b)** As the seismic waves passes, the vibrations moves the case (and coil) and causes the movement of the case (and coil) relative to the magnet. B1

This results in a change of magnetic flux linkage through the coil, and by Faraday's law, an e.m.f. will be induced across the coil. B1

**5(c)** 
$$\varepsilon = \frac{dN\phi}{dt} \Rightarrow \frac{d\phi}{dt} = \frac{\varepsilon}{N}$$

$$= \frac{85 \times 10^{-3}}{50}$$

$$= 1.7 \times 10^{-3}$$
 C1  
A1

**5(d)**   
 ✓ magnet with stronger magnetic field B1  
 ✓ coil with many turns B1  
 ✓ spring with a lower spring constant.  
 (any two of the above)

Question	Marks
6(a)	The r.m.s value of an <u>a.c.</u> will give the same <u>heating effect on a resistor</u> as due to <u>a d.c. current</u> of the <u>same value.</u> B1
6(b)(i)	Power = $VI = 1200 \text{ W}$ r.m.s current $I = 1200 / 120 = 10 \text{ A}$ Peak Current $I_o = 10\sqrt{2} = 14.1 \text{ A}$ C1 A1
6(b) (ii)	$I = I_o \sin (2\pi f t)$ $= 14.1 \sin (2\pi \times 0.50t)$ $= 14.1 \sin (100 \pi t)$ B1
6(c)	Secondary power = $600 + 1200 = 1800 \text{ W}$ By conservation of energy, Input power = $1800 \text{ W}$ C1 $I_p V_p = 1800$ $I_p = 1800 / (2.4 \times 10^3)$ $= 0.75 \text{ A}$ A1
7(a)(i)	an electric current that periodically reverses its direction in a circuit (with a frequency) B1
7(a)(ii)	- wire carries a current that is perpendicular to the magnetic field between the magnet poles B1 - (by Fleming's LHR) wire experiences a force that is normal to both the current and magnetic field B1 - force experienced is directed up and down vertically - since current reverses direction periodically, vibration is the same frequency as the AC B1
7(a)(iii)1.	recognising $\lambda = 2(L_{XP})$ M1 $v = f\lambda = (120)(4) = 480 \text{ m s}^{-1}$ A1
7(a) (iii)2.	- <u>meeting</u> of (waves) B1 - waves of same type, same frequency, opposite direction (characteristics) B1 - incident waves from force on AC-carrying wire in magnetic field reflected off X and P (Origin of standing waves) B1 - speed of incident / reflected wave in wire (concluding with correct answer) B1

Question	Marks
7(a) (iv) $v = f\lambda$ , therefore $v \propto f$ because $\lambda$ is fixed by length of XP tension is provided by weight, $T = mg$ therefore $T \propto m$	B1 B1
Need to show that $f \propto \sqrt{m}$ Take ratio:	
$\frac{f_1}{f_0} = \frac{170}{120} \approx 1.42$ , $\sqrt{\frac{2m}{m}} = \sqrt{2} \approx 1.41$	M1
$\frac{f_2}{f_0} = \frac{208}{120} \approx 1.73$ , $\sqrt{\frac{3m}{m}} = \sqrt{3} \approx 1.73$	
Likely valid	A0
7(b)(i) waves spread at edge or slit changes direction and spreads into geometric shadow	B1
7(b)(ii) $\theta = \frac{1}{2}(188 - 160) = 14^\circ$	C1
$d \sin \theta = n\lambda \rightarrow d \sin(14^\circ) = 2(633 \times 10^{-9})$	M1
$d = 5.23 \times 10^{-6}$ m	C1
$d = \frac{1}{N} \rightarrow N = \frac{1}{d} = 1.91 \times 10^5$	A1
7(b) (iii) each bright fringe is constructive interference from multiple points of diffracted waves (vs two points in double slit)	B1
<hr/>	
8(a) (i) loss in $E_k =$ gain in electric $E_p$	M1
$7.7 \times 10^{-13} = \frac{Qq}{4\pi\epsilon_0 r}$	
$= \frac{(79 \times 1.60 \times 10^{-19})(2 \times 1.60 \times 10^{-19})}{4\pi(8.85 \times 10^{-12})r}$	M1
$r = 4.72 \times 10^{-14}$ m	
Since $r$ is the distance of closest approach, the radius of gold must be less than this.	
8(a) (ii) 1. most of the atom is empty space or the size of the nucleus is very small compared to the size of the atom	B1
2. the nucleus is positively charged the mass is concentrated in the nucleus	B1 B1
8(b)(i) $7_{-1}^0\text{e}$	A1

Question		Marks
8(b)(ii)	mass defect = $[(82 \times 1.00863) + (57 \times 1.00728) - 138.955]u$ $= 1.16762u$ binding energy per nucleon $= \frac{1.16762 \times 1.66 \times 10^{-27} \times (3.0 \times 10^8)^2}{139 \times (1.60 \times 10^{-19})}$ $= 7.84 \text{ MeV}$	C1  M1  A1
8(b)(iii)	Since the fission reaction releases energy, the binding energy of uranium must be lower than for the products  <b>OR</b>  Above $A = 56$ , binding energy per nucleon decreases as $A$ increases. Since $^{235}\text{U}$ $^{235}\text{U}$ has a larger nucleon number (compared to $^{139}\text{La}$ ), it must have a higher binding energy per nucleon.	M1 A1  M1
8(c)(i)	<ul style="list-style-type: none"> <li>✓ radiation emitted in all directions</li> <li>✓ background radiation</li> <li>✓ emission from radioactive daughter products</li> <li>✓ window of detector may absorb some radiation</li> <li>✓ self-absorption in source</li> </ul>	Any 2  B1 each
8(c)(ii)	the energy is shared with a(n) (anti)neutrino	B1
8(c)(iii)	energy levels in the nuclei are discrete	B1
8(d)(i)	Background count = $10 \text{ min}^{-1}$ After $3 T_{\text{half}}$ , count rate will be $30 \text{ min}^{-1}$ Hence $3 T_{\text{half}} = 4.5 \text{ hours}$ $T_{\text{half}} = 1.5 \text{ hours (1.4 to 1.6 hours)}$	B1 M1  A1
8(d)(ii)	1. <u>no change</u> as radioactive <u>decay is spontaneous</u> / independent environment. 2. likely to be <u>different</u> as radioactive <u>decay is random</u> (and cannot be predicted).	B1 B1