

2

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the quadratic equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

3

- 1 A cylinder has a radius of $(\sqrt{2}-1)$ cm and a volume of $(12+3\sqrt{2})\pi$ cm³.

Find, **without using a calculator**, the exact value of its height, h cm, in the form $a+b\sqrt{2}$, where a and b are integers.

[3]

[Turn over

4

- 2 Suppose that x and y are non-zero real numbers such that $\frac{x}{3} = y^2$ and $\frac{x}{9} = 9y$.

Find the value of $x - y$.

[3]

5

- 3 Express $y = -2x^2 - 12x + 1$ in the form $y = a - 2(x + b)^2$ and hence state the maximum value of y , and its corresponding value of x . [4]

[Turn over

6

- 4 A curve is such that $\frac{dy}{dx} = \frac{2}{x^2} + \frac{3}{(7-2x)}$ and $P(3, 1)$ is a point on the curve.

Find the equation of the curve.

[4]

7

5 Express $\frac{2x-1}{x^2(x+1)}$ in partial fractions.

[6]

[Turn over

8

- 6 The function f is defined by $f(x) = 3x^3 - 5cx^2 + kc^2x + 4c^3$, where c and k are non-zero constants. $f(x)$ leaves a remainder of $-32c^3$ when divided by $x + 2c$.

(a) Find the value of k .

[2]

- (b) Using the value of k , determine whether $x^2 - cx - 2c^2$ is a factor of $f(x)$.
Justify your answer.

[3]

9

- 7 The function f is defined by $f(x) = 4 \cos ax + b$ for $0 \leq x \leq \pi$, where a and b are constants. The period of f is $\frac{2\pi}{3}$ and the function has a maximum value of 2.

(i) State the amplitude of f . [1]


(ii) Write down the value of a and of b . [2]

(iii) Sketch the graph of $y = f(x)$ for $0 \leq x \leq \pi$, indicating clearly the x -coordinates, in terms of π , of the points where the graph crosses the x -axis. [4]

[Turn over

10

- 8 In $\triangle ABC$, $\angle ABC = 90^\circ$, $AB = 12$ cm and $AC = 13$ cm. The rectangle $BPQR$ is such that its vertices P , Q and R lie on BC , CA and AB respectively.



It is given that $BP = x$ cm and $PQ = y$ cm.

(a) Show that $y = \frac{60 - 12x}{5}$.

[2]

(b) Show that the area, A cm², of the rectangle $BPQR$ is given by $A = 12x - \frac{12x^2}{5}$.

[1]


11

- 8 (c) Given that x can vary, find the stationary value of A and determine its nature. [4]

[Turn over

12

- 9 The diagram shows a quadrilateral $ABCD$ in which A is $(0, 6)$ and AB is parallel to the x -axis. D is a point on the x -axis such that the equation of DC is $x + 5y = -6$. AC is perpendicular to the line $2y - x = 7$.



(a) Find,

(i) the equation of AC ,

[2]

(ii) the coordinates of C .

[2]

13

- 9 (b) Given that the area of $\triangle ACD$ is 1.5 times that of $\triangle ABC$, find the coordinates of B .

[3]

- (c) Showing your working clearly, explain whether $ABCD$ is a kite.

[2]

[Turn over

14

10 (a) Prove the identity $\frac{1 - \sin^4 x}{\sin^2 x} = \cot^2 x + \cos^2 x$. [4]


15

- 10 (b) Hence solve the equation $\cot^2 2x + \cos^2 2x = 0$ for $0^\circ < x < 180^\circ$. [4]

[Turn over

16

- 11 The diagram shows a quadrilateral $ABCD$ whose vertices lie on the circumference of a circle. The point E lies on the extended line CD such that AE is a tangent to the circle at A . CD and AB are parallel lines.



- (a) Explain why angle $CBA =$ angle EDA . [2]

- (b) Show that triangles DAE and BAC are similar. [2]

17

- 11 (c) Given that $AD = DE$, explain why the line AC bisects the angle BCD . [3]

[Turn over

18

12 (a) Given that $2^{x-2} \times 3^{x+2} = 6^{2x}$, show that $x = \frac{\lg 9 - \lg 4}{\lg 6}$ [4]

19

12 (b) Solve $\log_9 y - 2 = \log_3 2y$.

[4]

[Turn over

20

- 13 (a) Water leaks from a container at a rate of $150 \text{ cm}^3/\text{s}$. The volume, $V \text{ cm}^3$, of the water in the container, when the height of water is $h \text{ cm}$, is given by

$$V = 10\pi + \frac{4\pi h^3}{9}. \text{ When } V = 334\pi \text{ cm}^3, \text{ find the}$$

- (i) value of h , [2]

- (ii) rate of change of h at this instant, correct to 3 significant figures. [3]

21


- 13 (b) The stopping distance, s m, of a car moving at v km/h can be modelled by the formula $\frac{s}{v} = \frac{1}{6} + \frac{v}{50}$.

(i) Find the rate at which s is changing with respect to v when $v = 60$. [3]

(ii) Explain the meaning of your answer to part (i). [1]

[Turn over

14



The diagram shows part of the curve $y = \sqrt{3x-2}$. The normal to the curve at B meets the x -axis at C . Given that the x -coordinate of B is 9, find

- (a) the coordinates of A and of C ,

[5]

23

14 (b) the area of the shaded region.

[5]

~ End of Paper ~

24

BLANK PAGE

2

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

3

- 1 By using an appropriate substitution, solve the equation $e^{\frac{1}{2}x} = 2 + 24e^{-\frac{1}{2}x}$. [5]

4

2 (a) Integrate $\left(\frac{6}{e^{3-x}}\right)^2$ with respect to x . [3]

(b) Explain, with working, why the equation $2x^3 + 3x^2 + 2x + 8 = 0$ has only 1 real root. Hence find this root. [5]


5

- 3 (a) Given that $y = x^2\sqrt{2x-1}$, show that $\frac{dy}{dx} = \frac{x(5x-2)}{\sqrt{2x-1}}$. Determine whether $y = x^2\sqrt{2x-1}$ is always an increasing function. [5]

- (b) Hence evaluate $\int_1^5 \frac{5x^2 - 2x + 1}{\sqrt{2x-1}} dx$. [4]

6

- 4 The diagram shows the front view of a rectangular block $ABCE$, with dimensions 14 cm by 8 cm, placed on a ramp, VE , tilted at an acute angle of θ° and $\angle VTE = 90^\circ$. The ramp is placed on a horizontal surface ET and d is the perpendicular distance from B to ET .



- (a) Show that $d = 14 \sin \theta + 8 \cos \theta$. [2]

- (b) Express d in the form $R \sin(\theta + \alpha)$, where $R > 0$ and $0^\circ \leq \alpha \leq 90^\circ$. [3]

7

(c) State the maximum value of d and find the corresponding value of θ . [2]

(d) Find the smallest value θ such that $d = 13$. [2]

8

- 5 (a) Explain by completing the square that $x^2 - 2x + 3$ is always positive for all real values of x . [2]

- (b) Hence, find the range of values of p if the inequality $\frac{3x^2 + px + 3}{x^2 - 2x + 3} > 0$ is satisfied for all real values of x . [4]

9

- 6 (a) In the binomial expansion of $\left(x + \frac{k}{x}\right)^5$, where k is a positive integer, the coefficients of $\frac{1}{x^3}$ and $\frac{1}{x}$ are the same. Find the value of k . [5]


- (b) Without expanding all the terms, explain why there is no constant term in the expansion of $(1 - 3x^2)\left(x + \frac{k}{x}\right)^5$. [3]

- 7 (a) The variables x and y , are related by the equation $\frac{3y^2}{k} + \frac{2x}{h} = 6$,
 where h and k are constants. When the graph of y^2 against x is plotted, a straight line is obtained. Given that the intercept on the y^2 axis is 8 and that the gradient of the line is -6 , calculate the value of h and of k . [3]

- (b) A glass of hot liquid is put on a table to cool. The temperature of the liquid, $T^\circ\text{C}$, after n minutes is given by $T = 25 + pe^{-kn}$, where p and k are constants. The table below shows the measured values of T and n .

n (minutes)	10	20	30	40	50
T ($^\circ\text{C}$)	79.6	58.1	45.1	37.2	32.4

- (i) Using the grid on page 11, plot $\ln(T - 25)$ against n and draw a straight line graph. [3]



12

Use your graph to estimate

(ii) value of p and of k , [4]

(iii) the temperature of the liquid after 35 minutes, [2]

13

- (iv) the number of minutes it takes for the temperature of the liquid to drop by 75% of its initial value. [2]

14

- 8 The velocity, $v \text{ ms}^{-1}$, of a particle, moving in a straight line, t seconds after motion has begun, is given by $v = 6t^2 + kt + 12$, where k is a constant. The particle passes a fixed point O with an acceleration of -6 ms^{-2} when $t = 1$.

(a) Show that $k = -18$. [2]


(b) Hence, find
(i) the minimum velocity achieved by the particle, [3]

15

- (ii) the total distance travelled during the first 4 seconds. [5]

16

9



- (a) The coordinates of the centre of a circle C_1 is $C(-5, 3)$. If the y -axis is a tangent of the circle, find the equation of the circle. [1]


A straight line cuts the circle at points A and B such that AB is parallel to $y + 3x - 11 = 0$ and that AC is parallel to the y -axis.

- (b) Find
- (i) the equation of the line AB and the equation of the perpendicular bisector of AB , [5]

(ii) hence, find the coordinates of B . [3]

(c) Circle C_2 is obtained by reflecting circle C_1 in the line AB . Find the equation of circle C_2 . [3]

- 10 The diagram shows part of the curve $y = 2 \sin 2x + 1$. P is the maximum point of the curve and Q is the point on the curve at which the gradient of the tangent is -4 .



- (a) Find the coordinates of P and of Q . [5]

19

- (b) Find the area of the shaded region bounded by the curve, the axes, line PQ and vertical line QR . Leave your answer in the exact form. [4]

-End of Paper-

BLANK PAGE