



**JUNYUAN SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2021
SECONDARY FOUR EXPRESS**

CANDIDATE NAME

CLASS

| | | |
|----------|----------|--|
| 4 | E | |
|----------|----------|--|

INDEX NUMBER

| | |
|--|--|
| | |
|--|--|

CHEMISTRY

6092

PAPER 1 Multiple Choice

xx September 2021

1 hour

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in.

Write in soft pencil.

Do not use paper clips, glue or correction fluid.

There are forty questions on this paper. Answer all questions. For each question there are four possible answers, A, B, C and D.

Choose the one you consider correct and record your choice in soft pencil on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

A copy of the Periodic Table is printed on page 18.

The use of an approved scientific calculator is expected, where appropriate.


This document consists of 18 printed pages.

2

- 1 Lead(II) iodide is insoluble in water.

Lead(II) iodide is prepared by adding aqueous lead(II) nitrate to potassium iodide.


Which pieces of apparatus are needed for the preparation and collection of solid lead(II) iodide from 20 cm³ of aqueous lead(II) nitrate?



- A 1, 2 and 4 B 1, 3 and 5 C 1, 4 and 5 D 2, 4 and 5

- 2 Some substances may be separated using paper chromatography. The diagram shows the results of running two mixtures in a suitable solvent.

Which spot has an R_f value of 0.38?



3

- 3 Compound X is a crystalline solid at room temperature and pressure. An aqueous solution of X is tested as shown.

| test | test result |
|---|---|
| acidify with dilute nitric acid, then add aqueous barium nitrate | no visible change |
| add aqueous ammonia | white precipitate, soluble in excess |

What could be the identity of X?

- A ammonium carbonate
 B sodium sulfate
 C calcium nitrate
 D zinc chloride
- 4 Gases are separated from liquid air by fractional distillation.

The boiling points of four gases are shown.

Which gas is both monatomic and a liquid at $-200\text{ }^{\circ}\text{C}$?


| | gas | boiling point / $^{\circ}\text{C}$ |
|---|----------|------------------------------------|
| A | argon | -186 |
| B | helium | -269 |
| C | neon | -246 |
| D | nitrogen | -196 |

4

- 5 Hydrogen chloride gas, HCl , reacts with ammonia gas, NH_3 , to form solid ammonium chloride.


The apparatus is set up as shown.

After a few minutes, solid ammonium chloride forms where the two gases meet.



The experiment is repeated using hydrogen bromide, HBr , in place of hydrogen chloride.

How far along the tube does the solid ammonium bromide form?



- 6 The table shows the numbers of particles present in the nuclei of four atoms or ions.

| | protons | neutrons | electronic structure |
|---|---------|----------|----------------------|
| 1 | 18 | 22 | 2.8.8 |
| 2 | 19 | 20 | 2.8.8 |
| 3 | 19 | 21 | 2.8.8.1 |
| 4 | 20 | 20 | 2.8.8.2 |

Which two particles belong to the same element?

- A** 1 and 2 **B** 1 and 4 **C** 2 and 3 **D** 2 and 4


- 7 Element X has a proton number of 20 and a nucleon number of 40. X reacts with element Y to form an ionic compound.

Which equation shows the process that takes place when X forms ions?


- A $X + e^- \rightarrow X^+$
 B $X + 2e^- \rightarrow X^{2+}$
 C $X \rightarrow X^+ + e^-$
 D $X \rightarrow X^{2+} + 2e^-$
- 8 A covalent molecule M contains a total of four shared electrons.

What is M?

- A ammonia, NH_3
 B hydrogen chloride, HCl
 C methane, CH_4
 D water, H_2O
- 9 Which pair of statements about diamond and graphite is correct?



diamond



graphite

- A Diamond and graphite are both pure carbon. They are both macromolecules.
 B Diamond and graphite can both be used as electrodes. Graphite is also used as a lubricant.
 C Diamond has covalent bonds. Graphite has ionic bonds.
 D Diamond is hard with a high melting point. Graphite is soft with a low melting point.
- 10 Which of the following contains the greatest mass of oxygen?
- A 0.2 mol of aluminium nitrate, $Al(NO_3)_3$
 B 0.3 mol of potassium sulfate, K_2SO_4
 C 0.4 mol of sodium nitrate, $NaNO_3$
 D 0.5 mol of magnesium carbonate, $MgCO_3$

6

- 11 Compound X has a composition by mass of 63.6 % nitrogen and 36.4 % oxygen.

What is the empirical formula of X?

- A N_2O B NO C NO_2 D N_2O_4

- 12 Calcium carbonate reacts with dilute hydrochloric acid according to the equation shown.



10 g of calcium carbonate is reacted with 100 cm³ of 1 mol/dm³ hydrochloric acid.


The following statements are made.

- 1 1.20 dm³ of carbon dioxide is formed.
- 2 5.55 g of calcium chloride is formed.
- 3 4.80 g of carbon dioxide is formed.
- 4 No calcium carbonate is left when the reaction is completed.

Which statements about the reaction are correct?

- A 1 and 2 B 1 and 4 C 2 and 3 D 3 and 4

- 13 The reaction for the conversion of bromoethane to ethanol is shown.



In an experiment, 10.90 g of bromoethane is converted to 3.45 g of ethanol.

What is the percentage yield of ethanol? [M_r : $\text{C}_2\text{H}_5\text{Br}$, 109; $\text{C}_2\text{H}_5\text{OH}$, 46]

- A 32 % B 42 % C 75 % D 100 %


- 14 Concentrated aqueous sodium chloride is electrolysed using inert electrodes.

Which row shows what is likely to happen in this electrolysis and why it happens?

| | change occurring | explanation |
|---|--|---|
| A | Oxygen is discharged at the anode. | $\text{OH}^{\ominus}(\text{aq})$ loses electrons more easily than does $\text{Cl}^{\ominus}(\text{aq})$. |
| B | During electrolysis, the pH of the electrolyte increases. | With the discharge of $\text{H}^{\oplus}(\text{aq})$ ions, $\text{OH}^{\ominus}(\text{aq})$ ions remain. |
| C | Solid sodium is discharged at the cathode. | $\text{Na}^{\oplus}(\text{aq})$ is present in aqueous solution. |
| D | The products stay the same if the aqueous sodium chloride is replaced by molten sodium chloride. | Na^{\oplus} and Cl^{\ominus} are present in both molten and aqueous sodium chloride. |

7

- 15 Pairs of metals are connected together to make a simple cell, as shown.



The table shows the reading on the voltmeter when different metals are used.


| | | metal 2 | | | |
|---------|-----------|-----------|---------|---------|-----------|
| | | beryllium | cerium | cobalt | manganese |
| metal 1 | beryllium | 0.00 V | +0.64 V | -1.57 V | -0.67 V |
| | cerium | | 0.00 V | -2.21 V | -1.30 V |
| | cobalt | | | 0.00 V | +0.90 V |
| | manganese | | | | 0.00 V |

If metal 2 is more reactive than metal 1, the voltage measured is positive.

What is the order of reactivity?

| | most reactive | → | | | least reactive |
|----------|---------------|-----------|-----------|-----------|----------------|
| A | cerium | beryllium | cobalt | manganese | |
| B | cerium | beryllium | manganese | cobalt | |
| C | cobalt | manganese | beryllium | cerium | |
| D | cobalt | manganese | cerium | beryllium | |


- 16 The energy level diagram for the reaction between P and Q to form R and S is shown.



Which row describes the energy changes involved and the type of reaction?

| | energy changes involved | type of reaction |
|----------|---|------------------|
| A | more energy is given out when the bonds in the products are formed than is absorbed to break the bonds in the reactants | endothermic |
| B | more energy is given out when the bonds in the products are formed than is absorbed to break the bonds in the reactants | exothermic |
| C | more energy is absorbed to break the bonds in the reactants than is given out when the bonds in the products are formed | endothermic |
| D | more energy is absorbed to break the bonds in the reactants than is given out when the bonds in the products are formed | exothermic |

- 17 Nitrogen trifluoride, NF_3 , is used in the manufacturing of certain types of solar panels. The equation for the formation of nitrogen trifluoride is shown.



| type of bond | bond energy (kJ/mol) |
|--------------------------|----------------------|
| $\text{N}\equiv\text{N}$ | +950 |
| $\text{F}-\text{F}$ | +150 |
| $\text{N}-\text{F}$ | +280 |

Using the table of bond energies, what is the energy change for this reaction?

- A** -560 kJ/mol
B -280 kJ/mol
C +280 kJ/mol
D +3080 kJ/mol

- 18 Marble chips react with hydrochloric acid in an exothermic reaction.



When excess marble chips are added to dilute hydrochloric acid, the rate of the reaction starts off fast, then gets slower until the reaction stops.

Why does the reaction rate get slower?


- A The concentration of the hydrochloric acid is decreasing.
 B The concentration of calcium chloride is increasing.
 C The calcium carbonate is completely used up.
 D The temperature of the mixture decreases.
- 19 Three experiments are carried out in which the same mass of magnesium is reacted with the same volume of dilute sulfuric acid at room temperature. The magnesium is in excess.

experiment 1 Large pieces of magnesium are used.

experiment 2 Small pieces of magnesium are used.

experiment 3 Large pieces of magnesium are used but the concentration of the acid is increased.


Graphs of the results are shown.



Which row is correct?

| | experiment 1 | experiment 2 | experiment 3 |
|---|--------------|--------------|--------------|
| A | W | X | Y |
| B | X | Y | W |
| C | Y | W | X |
| D | Y | X | W |

- 20 The reaction between magnesium and carbon dioxide is shown in the equation.



Which statement describes what happens in this reaction?

- A** Carbon is oxidised.
B Magnesium is reduced.
C Neither oxidation nor reduction happens.
D The carbon in carbon dioxide is reduced.
- 21 Gas X turns acidified potassium manganate(VII) from purple to colourless.
 Gas Y turns aqueous potassium iodide from colourless to brown.

What do these observations show about gas X and gas Y?

| | gas X | gas Y |
|----------|-----------------|-----------------|
| A | oxidising agent | oxidising agent |
| B | oxidising agent | reducing agent |
| C | reducing agent | oxidising agent |
| D | reducing agent | reducing agent |

- 22 An alloy of copper and zinc is added to an excess of dilute hydrochloric acid. The resulting mixture is then filtered.

Which observations are correct?

| | filtrate | residue |
|----------|---------------------|-----------|
| A | colourless solution | none |
| B | colourless solution | red-brown |
| C | blue solution | grey |
| D | blue solution | non |

- 23 Three chemicals, P, Q and R, were each dissolved in water. The table shows some of the reactions of these solutions.


| solution | reaction when solid sodium carbonate is added | reaction when heated with solid ammonium chloride |
|----------|---|---|
| P | gas evolved | no reaction |
| Q | no reaction | gas evolved |
| R | no reaction | no reaction |

The pH of the three solutions was also measured.

What are the likely pH values of these solutions?

| | P | Q | R |
|---|----|----|----|
| A | 2 | 7 | 13 |
| B | 2 | 13 | 7 |
| C | 7 | 2 | 13 |
| D | 13 | 7 | 2 |

- 24 The diagram shows the steps in the preparation of a salt.



Which salt can be prepared by this method?

- A barium sulfate
- B copper(II) sulfate
- C potassium sulfate
- D sodium sulfate

- 25 Three different elements react by losing electrons. The ions formed all have the electronic configuration, 2.8.

Which statement about these elements is correct?

- A They are noble gases.
 B They are transition elements.
 C They are in the same group.
 D They are in the same period.
- 26 Sodium is a Group I metal.
- Which property, that is typical of most metals, is not shown by sodium?
- A conductor of electricity
 B high melting point
 C malleable
 D shiny
- 27 Palladium is an element with the atomic number of 46. Some of its properties, and the properties of its compounds, can be predicted from its position in the Periodic Table.

Which row is correct?

| | predicted property of palladium | predicted property of palladium compounds |
|----------|--|---|
| A | Its density is similar to the density of sodium. | Some of them can act as catalysts. |
| B | Its density is similar to the density of sodium. | They are white in the solid state. |
| C | It is present in compounds in more than one oxidation state. | Some of them can act as catalysts. |
| D | It is present in compounds in more than one oxidation state. | They are white in the solid state. |

- 28 The metal beryllium does not react with cold water.

It reacts with hydrochloric acid but cannot be extracted from its ore by using carbon.

Where is beryllium placed in the reactivity series?

magnesium

A

zinc

B


iron

C

copper

D


- 29 Four equations are shown.



Which equations represent reactions that take place in the blast furnace where iron is extracted from haematite?

- A** 1, 2 and 3 **B** 1, 2 and 4 **C** 2, 3 and 4 **D** 3 and 4 only

- 30 Methanol is made by reacting carbon monoxide with hydrogen. The reaction is reversible.



The forward reaction is exothermic.

What combination of temperature and pressure gives the highest equilibrium yield of methanol?

| | temperature / °C | pressure / atmospheres |
|----------|---------------------|---------------------------|
| A | 200 | 10 |
| B | 200 | 200 |
| C | 500 | 10 |
| D | 600 | 200 |

- 31 Which pair of gases could be removed from the atmosphere using calcium carbonate?
- A** CO₂ and O₃
B CO and SO₂
C CH₄ and NO₂
D NO₂ and SO₂

- 32 Oxides of nitrogen, such as NO and NO₂, are formed in the petrol engines of cars.


They are removed from the exhaust gases by reactions in the car's catalytic converter.

Which row describes how oxides of nitrogen are formed in a petrol engine, and a reaction that happens in the catalytic converter?

| | how oxides of nitrogen are formed | a reaction that happens in the catalytic converter |
|----------|--|---|
| A | by the reaction between nitrogen and oxygen from the air | $2\text{NO} + 2\text{CO} \rightarrow \text{N}_2 + 2\text{CO}_2$ |
| B | by the reaction between nitrogen and oxygen from the air | $2\text{NO} + 2\text{H}_2 \rightarrow \text{N}_2 + 2\text{H}_2\text{O}$ |
| C | by the reaction between nitrogen compounds in petrol and oxygen from the air | $2\text{NO} + 2\text{CO} \rightarrow \text{N}_2 + 2\text{CO}_2$ |
| D | by the reaction between nitrogen compounds in petrol and oxygen from the air | $2\text{NO} + 2\text{H}_2 \rightarrow \text{N}_2 + 2\text{H}_2\text{O}$ |

- 33 The pie chart represents the composition of natural gas.

Which sector represents methane?




- 34 When a molecule of a saturated hydrocarbon is cracked, it forms two molecules, X and Y.

Which row is correct?

| | X | Y |
|----------|------------------|----------------------------------|
| A | H ₂ O | C _n H _{2n} |
| B | H ₂ O | C _n H _{2n+2} |
| C | H ₂ | C _n H _{2n} |
| D | H ₂ | C _n H _{2n+2} |

- 35 Which diagram shows a branched-chain isomer of butane?



- 36 Which organic compound requires the least number of moles of oxygen for the complete combustion of one mole of the compound?

A C₃H₇OH **B** C₃H₇COOH **C** C₃H₈ **D** C₄H₈


- 37 Chlorine reacts with methane.

Which row is correct?

| | chemical equation | conditions required |
|----------|--|---|
| A | $Cl_2 + CH_4 \rightarrow CH_2Cl_2 + H_2$ | methane and chlorine gases are mixed in the presence of ultraviolet light |
| B | $Cl_2 + CH_4 \rightarrow CH_2Cl_2 + H_2$ | methane is bubbled into concentrated aqueous chlorine |
| C | $Cl_2 + CH_4 \rightarrow CH_3Cl + HCl$ | methane and chlorine gases are mixed in the presence of ultraviolet light |
| D | $Cl_2 + CH_4 \rightarrow CH_3Cl + HCl$ | methane is bubbled into concentrated aqueous chlorine |





- 38 Fats are essential components of the human diet.

The diagram shows a fat molecule.







Which description of this fat molecule is correct?

- A saturated carboxylic acid
 B saturated ester
 C unsaturated carboxylic acid
 D unsaturated ester
- 39 Which partial structure is correct for the product of polymerization of butane, $\text{CH}_2=\text{CHCH}_2\text{CH}_3$?



40 Ethanoic acid is reacted with propanol.

What is the name and structure of the ester produced?

| | name | structure |
|----------|------------------|--|
| A | propyl ethanoate |  |
| B | ethyl propanoate |  |
| C | propyl ethanoate |  |
| D | ethyl propanoate |  |

End of Paper

The Periodic Table of Elements

| | | Group | | | | | | | | | | | | | | | |
|----------------------------|-----------------------------|-----------------------------|---------------------------------|-----------------------------|------------------------------|------------------------------|------------------------------|------------------------------|--------------------------------|-------------------------------|-------------------------------|-----------------------------|-------------------------------|-----------------------------|-------------------------------|-------------------------------|------------------------------|
| I | II | III | IV | V | VI | VII | 0 | | | | | 0 | | | | | |
| 3 Li lithium 7 | 4 Be beryllium 9 | 5 B boron 11 | 6 C carbon 12 | 7 N nitrogen 14 | 8 O oxygen 16 | 9 F fluorine 19 | 10 Ne neon 20 | | | | | 18 Ar argon 40 | | | | | |
| 11 Na sodium 23 | 12 Mg magnesium 24 | 13 Al aluminium 27 | 14 Si silicon 28 | 15 P phosphorus 31 | 16 S sulfur 32 | 17 Cl chlorine 35.5 | 18 Ar argon 40 | | | | | 36 Kr krypton 84 | | | | | |
| 19 K potassium 39 | 20 Ca calcium 40 | 21 Sc scandium 45 | 22 Ti titanium 48 | 23 V vanadium 51 | 24 Cr chromium 52 | 25 Mn manganese 55 | 26 Fe iron 56 | 27 Co cobalt 59 | 28 Ni nickel 58 | 29 Cu copper 64 | 30 Zn zinc 65 | 31 Ga gallium 70 | 32 Ge germanium 73 | 33 As arsenic 75 | 34 Se selenium 79 | 35 Br bromine 80 | 36 Kr krypton 84 |
| 37 Rb rubidium 85 | 38 Sr strontium 88 | 39 Y yttrium 89 | 40 Zr zirconium 91 | 41 Nb niobium 93 | 42 Mo molybdenum 96 | 43 Tc technetium - | 44 Ru ruthenium 101 | 45 Rh rhodium 103 | 46 Pd palladium 106 | 47 Ag silver 108 | 48 Cd cadmium 112 | 49 In indium 115 | 50 Sn tin 119 | 51 Sb antimony 122 | 52 Te tellurium 128 | 53 I iodine 127 | 54 Xe xenon 131 |
| 55 Cs caesium 133 | 56 Ba barium 137 | 57-71 lanthanoids | 72 Hf hafnium 178 | 73 Ta tantalum 181 | 74 W tungsten 184 | 75 Re rhenium 186 | 76 Os osmium 190 | 77 Ir iridium 192 | 78 Pt platinum 195 | 79 Au gold 197 | 80 Hg mercury 201 | 81 Tl thallium 204 | 82 Pb lead 207 | 83 Bi bismuth 209 | 84 Po polonium - | 85 At astatine - | 86 Rn radon - |
| 87 Fr francium - | 88 Ra radium - | 89-103 actinoids | 104 Rf rutherfordium - | 105 Db dubnium - | 106 Sg seaborgium - | 107 Bh bohrium - | 108 Hs hassium - | 109 Mt meitnerium - | 110 Ds darmstadtium - | 111 Rg roentgenium - | 112 Cn copernicium - | 114 Fl flerovium - | 116 Lv livermorium - | 118 Og oganeson - | 119 Uue unbinilium - | 120 Uub unbinilium - | 121 Uut ununilium - |

1 H
hydrogen
1

Key
proton (atomic) number
atomic symbol
name
relative atomic mass

lanthanoids

actinoids

| | | | | | | | | | | | | | | |
|------------------------------|----------------------------|---------------------------------|------------------------------|-----------------------------|-----------------------------|-----------------------------|-------------------------------|----------------------------|-------------------------------|------------------------------|---------------------------|-------------------------------|------------------------------|------------------------------|
| 57 La lanthanum 138 | 58 Ce cerium 140 | 59 Pr praseodymium 141 | 60 Nd neodymium 144 | 61 Pm promethium - | 62 Sm samarium 150 | 63 Eu europium 152 | 64 Gd gadolinium 157 | 65 Tb terbium 159 | 66 Dy dysprosium 163 | 67 Ho holmium 165 | 68 Er erbium 167 | 69 Tm thulium 169 | 70 Yb ytterbium 173 | 71 Lu lutetium 175 |
| 89 Ac actinium - | 90 Th thorium 232 | 91 Pa protactinium 231 | 92 U uranium 238 | 93 Np neptunium - | 94 Pu plutonium - | 95 Am americium - | 96 Cm curium - | 97 Bk berkelium - | 98 Cf californium - | 99 Es einsteinium - | 100 Fm fermium - | 101 Md mendelevium - | 102 No nobelium - | 103 Lr lawrencium - |

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).



JUNYUAN SECONDARY SCHOOL
Secondary Four Express
Preliminary Examination 2021

CANDIDATE
NAME

CLASS

INDEX
NUMBER

CHEMISTRY

6092 / 02

Paper 2

23 Aug 2021

1 hour 45 min

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, index number and name on all the work you hand in.
 Write in dark blue or black pen on both sides of the paper.
 You may use a soft pencil for any diagrams, graphs or rough workings.
 Do not use staples, paper clips, highlighters, glue or correction fluid.

The use of an approved scientific calculator is expected, where appropriate.

Section A

Answer **all** the questions in the spaces provided.

Section B

Answer all **three** questions, the last question is in the form either/or.
 Write your answers in the spaces provided.

At the end of the examination, fasten all your work securely together.
 The number of marks is given in brackets [] at the end of each question or part question.
 A copy of the Periodic Table is printed in page 21.

| For Examiner's Use | |
|--------------------|----|
| Section A | |
| Section B | |
| Total | 80 |


This document consists of **21** printed pages and 1 blank page.

2

Section A

Answer all questions in this section in the spaces provided.
The total mark for this section is 50.

A1 The bonding in compound YZX_2 , formed between atoms Y, Z, and X, is shown below.



(a) State the Groups that elements X and Z belong to in the Periodic Table.

Element X:[1]

Element Z:[1]

(b) Element Z has the smallest atomic radius in its Group. Identify element Z.

.....[1]

(c) Element Y has 16 protons. State the period in which element Y is found in the Periodic Table.

.....[1]

(d) Aluminium forms a compound with element X. Write the chemical formula of this compound.

.....[1]

(e) Another element, V, has 1 more proton than element X. State a property that element V is likely to have.

.....

.....[1]

[Total: 6]

3

- A2 (a)** In order to make computer processing easier, a system using proton (atomic) numbers rather than names or formulae has been developed. The following examples illustrate this code system.

| name | formula | code |
|------------------|--------------------------------|-------------|
| sodium chloride | NaCl | 11,17 |
| iron(II) bromide | FeBr ₂ | 26, 35(2) |
| aluminium oxide | Al ₂ O ₃ | 13(2), 8(3) |

Use this system to complete the table below.

| name | formula | code |
|------------------|---------|-----------|
| | KF | |
| | | 29,6,8(3) |
| nitrogen dioxide | | |

[3]

- (b)** A sample of chlorine gas was found to consist of isotopes Cl-35 and Cl-37. A calculation of the relative atomic mass of chlorine was found to be 36.5.

- (i)** Define isotopes.

.....[1]

- (ii)** For the two chlorine isotopes, compare and contrast:

- the relative abundance,
- and the atomic structure.

.....

 [3]

[7 marks]


4

A3 Concrete is an excellent building material that is commonly used in Singapore. It is made from cement, sand and water. Concrete is slightly porous. When rainwater soaks through concrete, some of the uncombined calcium oxide in the cement, dissolves to form aqueous calcium hydroxide.

(a) Write a balanced chemical equation, with state symbols for this reaction.

.....[2]

The diagram below shows the cross-section of a cracked concrete beam after a rainfall. Over time, air diffuses through the cracks to create a zone of reaction.



(b) Explain why the pH in a wet concrete beam is high.


.....
[1]

(c) Explain why there will be a chemical reaction when air diffuses into a wet concrete beam.

.....

[2]

To strengthen concrete and prevent cracks, steel reinforcement is often used as seen in the diagram below.



Source: <http://www.ashireporter.org/HomeInspection/Articles/Commercial-Inspection-CONCRETE-101/1068#crack-pattern.gif>

- (c) In terms of its structure, explain why steel is able to act as a reinforcement against compressive forces.

.....


[2]

- (d) Some modern structures, especially those situated near the sea, use fiberglass reinforcement instead of steel. Why is this so?

.....
[1]

[Total: 8]

- A4** A student wishes to identify the anions that are present in a sample of wastewater. The following flowchart shows how he can identify the anions that are present:



- (a) For anions A and B, write ionic equations for the formation of white precipitates described above. (Use the actual chemical symbols of the ions involved.)

.....
[2]

- (b) A student noticed that the test for the anion, nitrate, has been missed out in the flowchart above. He suggests that after observing if there is effervescence (i.e step 1), the resulting mixture should be divided into three samples to include a test for nitrate.

Explain why this will not give conclusive results.

.....

[2]


7

- (c) The wastewater contains only aluminium and calcium cations. Draw a flowchart that would enable him to confirm the identity of both ions, using aqueous sodium hydroxide and aqueous ammonia.


[3]

[Total: 7]

- A5** Hydrogen has many industrial uses. One possible way to manufacture hydrogen involves the reaction between methane and steam.



The diagram shows the energy profile diagram for this reaction.



- (a) State one industrial use of hydrogen.

.....[1]

- (b) What do the arrows, X and Y, represent in the diagram?

X:

Y:[2]

6092/02/O/4E/Prelim/21

8

- (c) A student wrote the following statement:

"In the reaction between methane and steam, the amount of energy released for bond forming is smaller than the amount of energy absorbed for bond breaking."

Do you agree with the student? Explain your answer with reference to the energy profile diagram.

.....


 [2]

- (d) If a catalyst can be found for this reaction, draw on the diagram above, how the energy profile will be change.

[1]

[Total: 6]

- A6 A student set up two experiments for electroplating steel keychains with copper. She closed both circuits for a period of time.



- (a) Complete the table of information about the experiments.

| experiment | electrodes | ionic equation, for reaction at each electrode |
|------------|----------------|--|
| 1 | carbon rod | |
| | steel keychain | |
| 2 | copper rod | |
| | steel keychain | |

[3]

(b) Describe and explain the observations seen in the electrolyte in each experiment.

Experiment 1:

.....
.....
.....

Experiment 2:

.....
.....
.....[4]

(c) The student repeated Experiment 2 with another keychain. However, she left the keychain in the setup without closing the switch for days.

Describe and explain one change she would observe of the keychain.

.....
.....
.....
.....[3]

[Total: 10]

10

A7 Ketones are a homologous series of organic compounds.

The table shows the names, formulae and boiling points of some ketones.

| name | structural formula | boiling point / °C |
|-------------|---|--------------------|
| 2-propanone | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{C}-\text{CH}_3 \end{array}$ | 56 |
| 2-butanone | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{C}-\text{CH}_2-\text{CH}_3 \end{array}$ | 80 |
| 3-pentanone | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{CH}_2-\text{C}-\text{CH}_2-\text{CH}_3 \end{array}$ | |
| 3-hexanone | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3\text{CH}_2-\text{C}-\text{CH}_2\text{CH}_2\text{CH}_3 \end{array}$ | 123 |

(a) Deduce the general formula and functional group of the ketone homologous series.

general formula functional group [2]


(b) Predict the boiling point of 3-pentanone.

.....[1]


(c) Using the information above, deduce the significance of the number "3" in the name "3-hexanone".

.....[1]

(d) 2-propanone, propanal (an aldehyde) and 2-propanol are all compounds of carbon, hydrogen and oxygen.



propanal



2-propanol

State with reasons, which of these three compounds are isomers of each other.

.....

[2]

[Total: 6]

6092/02/O/4E/Prelim/21

SECTION B

Answer all three questions from this section.

The last question is in the form of either/or and only one of the alternatives should be attempted.
The total mark for this section is 30.

- B8** Hydrogen peroxide is a common chemical found in the laboratories and in solutions used in our daily lives. The following sources of information illustrate how hydrogen peroxide can be used in managing the spread of the coronavirus and also as a bleaching agent.

Source 1:

HYDROGEN PEROXIDE

HANDS **HARD SURFACES**

H_2O_2

HYDROGEN PEROXIDE

Don't mix peroxide with vinegar.
This makes corrosive peracetic acid.

MINIMUM CONCENTRATION OF 0.5% PEROXIDE

HOW DOES IT DESTROY THE VIRUS?

Peroxide oxidises and destroys virus proteins and genetic material. It should be left on surfaces for at least 10 minutes.

It is common to express the concentration of chemicals in a cleaning solution by volume %, e.g. a 50% hydrogen peroxide is formed by mixing 50 cm³ of hydrogen peroxide with 50 cm³ of water.

'ball-and-stick' model for the structure of peracetic acid

Adapted from info source:


<https://www.compoundchem.com/2020/03/31/destroy-coronavirus/>

6092/02/O/4E/Prelim/21

Source 2:

Hydrogen peroxide is often found in bleaching solutions as it exhibits strong bleaching properties upon its decomposition into oxygen gas and water. A compound called TEAD is then added to the bleach solution to improve bleaching performance.

The graph below shows the performance of a hydrogen peroxide containing bleach solution, at different temperatures, with and without TEAD.



- (a) Draw a "dot-and-cross" diagram to show the bonding in hydrogen peroxide. You are required to show only the outermost electrons.

[2]

13

(b) A bottle of hydrogen peroxide solution of concentration 1.0 mol/dm^3 is found in the lab.

(i) Find the concentration of the hydrogen peroxide in g/dm^3 . [Mr of $\text{H}_2\text{O}_2 = 34$]

[1]

(ii) Using your answer in b(i) and given that the density of hydrogen peroxide liquid is 1.45 g/cm^3 , show, using calculations, whether the bottle of hydrogen peroxide solution meets the minimum concentration required for disinfecting surfaces.

[2]

(c) Based on the structure of peracetic acid, predict two properties that it will have.

.....
.....
.....[2]

(d) Describe the trend in the bleaching performance between the bleaches using TEAD and without using TEAD with respect to temperature.

.....
.....
.....
.....[2]

14


- (e) The graph below shows the distribution of energy of the particles in the bleach at a given temperature T_1 .

Unshaded portion below the curve:

- represents colliding particles with energy less than E_a .

Shaded portion below the curve:

- represents colliding particles with energy equal or more than E_a .



- (i) On the diagram itself, draw another vertical dotted line to mark the new E_a when a suitable catalyst is added and the temperature remains at T_1 . Label the new E_a as $E_a(\text{catalyst})$. [1]
- (ii) Shade the new portion below the curve that represent effective collisions among particles after the catalyst is added. [1]
- (iii) What does your answer in (e)(ii) suggest about the effect of adding a catalyst?

.....
[1]

[Total: 12]

6092/02/O/4E/Prelim/21

B9 Mercury, Venus and Earth are the three inner planets in the solar system with Mercury closest to the sun. Venus is both the closest planet to Earth and the planet closest in size to Earth. With an average surface temperature of 462 °C, Venus is by far the hottest planet in the Solar System. Table 9.1 below shows the atmospheric composition on Venus.




Table 9.1: percentage composition of the atmosphere on Venus

| gas | percentage composition |
|-----------------|------------------------|
| carbon dioxide | 96.5 |
| nitrogen | 3.5 |
| sulfur dioxide | 0.0150 |
| argon | 0.0070 |
| water vapour | 0.0020 |
| carbon monoxide | 0.0017 |
| helium | 0.0012 |
| neon | 0.0007 |

(a) Describe how the percentage composition of the atmosphere on Venus differs from that on Earth, with respect to carbon dioxide and nitrogen.

.....

[2]

(b) Suggest why Venus has a much higher temperature than Mercury despite being further away from the Sun.

.....

[2]

- (d) Suggest one property of metal that need to be taken into consideration when using it to make a space probe to land on Venus. Explain your answer.

.....
.....
.....[2]

- (e) Describe a test to show that the atmosphere on Venus contains sulfur dioxide.


.....
.....
.....[2]

[Total: 8]


EITHER

B10 A student conducted an experiment to determine the molar mass of an unknown metal, X. The student mixes solid iodine with an excess of metal X. When water is added to the mixture, a reaction occurs between iodine and the metal to form a water-soluble compound, XI_2 .

The equation can be represented below:



The reaction proceeds until all of the iodine is used up. The XI_2 solution is collected and heated to remove the water and the product is dried and weighed to a constant mass. The experimental steps are represented in the diagram and the results are tabulated below.



| | |
|---|--------|
| mass of beaker / g | 125.46 |
| mass of beaker + metal X / g | 126.55 |
| mass of beaker + metal X + I_2 / g | 127.57 |
| mass of XI_2 only (first weighing) / g | 1.28 |
| mass of XI_2 only (second weighing) / g | 1.28 |

(a) Calculate the number of moles of I_2 used.

[2]

18

- (b) Calculate, to the nearest whole number, the molar mass of the unknown metal, X. Identify metal X.

Molar mass of X is

Metal X is[4]

- (c) The student hypothesized that

“the compound formed in the reaction, XI_2 , is ionic.”

Briefly suggest how the student could test out this hypothesis in the laboratory.

.....

[2]

- (d) The student also predicted that chlorine gas will react with metal X more vigorously than solid iodine at room temperature.

With reference to the trends observed in the Periodic Table, justify if this prediction would be true.

.....


[2]

[Total: 10]


6092/02/O/4E/Prelim/21

OR

B10 Hot glue, a form of thermoplastic adhesive in solid cylindrical sticks are melted in the electric glue gun with a continuous-duty heating element. The glue is sticky when hot, and solidifies within a minute.




The glue is a polymer known as ethylene vinyl acetate (also known as EVA). EVA is a polymer made up of two different polymers joined together. It is a copolymer of poly(ethylene) and poly(vinyl acetate). The structural formula of EVA is as shown below.




(a) Draw the full structural formula of the monomer used to form

- (i) poly(ethylene),
- (ii) poly(vinyl acetate).



monomer of poly(ethylene)



monomer of poly(vinyl acetate)

[2]

20

- (b) A sample of EVA polymer made with 20% by mass of vinyl acetate was analysed and found to have an average relative molecular mass of 12040.


How many carbon atoms are present in an average chain?

Show your working clearly.

[4]

- (c) (i) Two organic molecules, **M** and **N** can be used to synthesize the monomer of poly(vinyl acetate).

Draw the full structural formula of these organic molecules, **M** and **N**.



[2]

- (ii) State the conditions needed for the reaction between organic molecules, **M** and **N** to occur.

.....[1]

- (iii) Zinc granules were added into two beakers, each containing one of the molecules, **M** and **N**. Only one of the beakers showed a reaction.

Describe what was observed for the beaker which showed a reaction when zinc was added.

.....
[1]

[Total: 10]

6092/02/O/4E/Prelim/21

PartnerInLearning

437

More papers at www.testpapersfree.com

The Periodic Table of Elements

| I | | II | | Group | | | | | | | | | | III | IV | V | VI | VII | 0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|-----------------------------|---------------------------|-----------------------------|-----------------------------|--|---------------------------|----------------------------|----------------------------|---------------------------|----------------------------|-----------------------------|------------------------|--------------------------|--------------------------|-----------------------------|---------------------------|-----------------------------|-----------------------------|------------------------------|----------------------------|---------------------------|----------------------------|----------------------------|-----------------------------|----------------------------|-----------------------------|---------------------------|----------------------------|---------------------------|--------------------------|------------------------|----------------------------|-----------------------------|----------------------------|----------------------------|---|----------------------------|-----------------------------|-------------------------|----------------------------|-----------------------------|-------------------------|----------------------------|---------------------------|---------------------------|-----------------------------|---------------------------|-----------------------------|-------------------------|------------------------------|-------------------------|---------------------------|----------------------------|----------------------------|---------------------------|----------------------------|-----------------------------|------------------------|--------------------------|--------------------------|--------------------------|------------------------|---------------------------|-----------------------------|---------------------------|----------------------------|---------------------------|---------------------------|------------------------|-----------------------------|------------------------------|--------------------------|--------------------------|
| 3 Li lithium 7 | 4 Be beryllium 9 | 11 Na sodium 23 | 12 Mg magnesium 24 | 19 K potassium 39 | 20 Ca calcium 40 | 21 Sc scandium 45 | 22 Ti titanium 48 | 23 V vanadium 51 | 24 Cr chromium 52 | 25 Mn manganese 55 | 26 Fe iron 56 | 27 Co cobalt 59 | 28 Ni nickel 59 | 29 Cu copper 64 | 30 Zn zinc 65 | 31 Ga gallium 70 | 32 Ge germanium 73 | 33 As arsenic 75 | 34 Se selenium 79 | 35 Br bromine 80 | 36 Kr krypton 84 | 37 Rb rubidium 85 | 38 Sr strontium 88 | 39 Y yttrium 89 | 57-71 lanthanoids | 56 Ba barium 137 | 55 Cs caesium 133 | 87 Fr francium — | 88 Ra radium — | 89-103 actinoids | 72 Hf hafnium 178 | 73 Ta tantalum 181 | 74 W tungsten 184 | 75 Re rhenium 186 | 76 Os osmium 190 | 77 Ir iridium 192 | 78 Pt platinum 195 | 79 Au gold 197 | 80 Hg mercury 201 | 81 Tl thallium 204 | 82 Pb lead 207 | 83 Bi bismuth 209 | 84 Po polonium — | 85 At astatine — | 86 Rn radon — | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | <table border="1"> <tr> <td>1 H hydrogen 1</td> <td colspan="10"></td> <td>5 B boron 11</td> <td>6 C carbon 12</td> <td>7 N nitrogen 14</td> <td>8 O oxygen 16</td> <td>9 F fluorine 19</td> <td>10 Ne neon 20</td> </tr> <tr> <td colspan="4"></td> <td colspan="10"> <table border="1"> <tr> <td>13 Al aluminium 27</td> <td>14 Si silicon 28</td> <td>15 P phosphorus 31</td> <td>16 S sulfur 32</td> <td>17 Cl chlorine 35.5</td> <td>18 Ar argon 40</td> </tr> </table> </td> <td>20 Ca calcium 40</td> <td>21 Sc scandium 45</td> <td>22 Ti titanium 48</td> <td>23 V vanadium 51</td> <td>24 Cr chromium 52</td> <td>25 Mn manganese 55</td> <td>26 Fe iron 56</td> <td>27 Co cobalt 59</td> <td>28 Ni nickel 59</td> <td>29 Cu copper 64</td> <td>30 Zn zinc 65</td> <td>31 Ga gallium 70</td> <td>32 Ge germanium 73</td> <td>33 As arsenic 75</td> <td>34 Se selenium 79</td> <td>35 Br bromine 80</td> <td>36 Kr krypton 84</td> </tr> </table> | | | | | | | | | | 1 H hydrogen 1 | | | | | | | | | | | 5 B boron 11 | 6 C carbon 12 | 7 N nitrogen 14 | 8 O oxygen 16 | 9 F fluorine 19 | 10 Ne neon 20 | | | | | <table border="1"> <tr> <td>13 Al aluminium 27</td> <td>14 Si silicon 28</td> <td>15 P phosphorus 31</td> <td>16 S sulfur 32</td> <td>17 Cl chlorine 35.5</td> <td>18 Ar argon 40</td> </tr> </table> | | | | | | | | | | 13 Al aluminium 27 | 14 Si silicon 28 | 15 P phosphorus 31 | 16 S sulfur 32 | 17 Cl chlorine 35.5 | 18 Ar argon 40 | 20 Ca calcium 40 | 21 Sc scandium 45 | 22 Ti titanium 48 | 23 V vanadium 51 | 24 Cr chromium 52 | 25 Mn manganese 55 | 26 Fe iron 56 | 27 Co cobalt 59 | 28 Ni nickel 59 | 29 Cu copper 64 | 30 Zn zinc 65 | 31 Ga gallium 70 | 32 Ge germanium 73 | 33 As arsenic 75 | 34 Se selenium 79 | 35 Br bromine 80 | 36 Kr krypton 84 | 50 Sn tin 119 | 51 Sb antimony 122 | 52 Te tellurium 128 | 53 I iodine 127 | 54 Xe xenon 131 |
| 1 H hydrogen 1 | | | | | | | | | | | 5 B boron 11 | 6 C carbon 12 | 7 N nitrogen 14 | 8 O oxygen 16 | 9 F fluorine 19 | 10 Ne neon 20 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | <table border="1"> <tr> <td>13 Al aluminium 27</td> <td>14 Si silicon 28</td> <td>15 P phosphorus 31</td> <td>16 S sulfur 32</td> <td>17 Cl chlorine 35.5</td> <td>18 Ar argon 40</td> </tr> </table> | | | | | | | | | | 13 Al aluminium 27 | 14 Si silicon 28 | 15 P phosphorus 31 | 16 S sulfur 32 | 17 Cl chlorine 35.5 | 18 Ar argon 40 | 20 Ca calcium 40 | 21 Sc scandium 45 | 22 Ti titanium 48 | 23 V vanadium 51 | 24 Cr chromium 52 | 25 Mn manganese 55 | 26 Fe iron 56 | 27 Co cobalt 59 | 28 Ni nickel 59 | 29 Cu copper 64 | 30 Zn zinc 65 | 31 Ga gallium 70 | 32 Ge germanium 73 | 33 As arsenic 75 | 34 Se selenium 79 | 35 Br bromine 80 | 36 Kr krypton 84 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 13 Al aluminium 27 | 14 Si silicon 28 | 15 P phosphorus 31 | 16 S sulfur 32 | 17 Cl chlorine 35.5 | 18 Ar argon 40 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Key
proton (atomic) number
atomic symbol
name
relative atomic mass

| | | | | | | | | | | | | | | |
|------------------------------|----------------------------|---------------------------------|------------------------------|-----------------------------|-----------------------------|-----------------------------|-------------------------------|----------------------------|-------------------------------|------------------------------|---------------------------|-------------------------------|------------------------------|------------------------------|
| 57 La lanthanum 139 | 58 Ce cerium 140 | 59 Pr praseodymium 141 | 60 Nd neodymium 144 | 61 Pm promethium — | 62 Sm samarium 150 | 63 Eu europium 152 | 64 Gd gadolinium 157 | 65 Tb terbium 159 | 66 Dy dysprosium 163 | 67 Ho holmium 165 | 68 Er erbium 167 | 69 Tm thulium 169 | 70 Yb ytterbium 173 | 71 Lu lutetium 175 |
| 89 Ac actinium — | 90 Th thorium 232 | 91 Pa protactinium — | 92 U uranium 238 | 93 Np neptunium — | 94 Pu plutonium — | 95 Am americium — | 96 Cm curium — | 97 Bk berkelium — | 98 Cf californium — | 99 Es einsteinium — | 100 Fm fermium — | 101 Md mendelevium — | 102 No nobelium — | 103 Lr lawrencium — |

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

http://www.wisesas.com/Chemistry/Organic/Naming_Ethers.ppt

6092/02/O/4E/Prelim/21

6092 PURE CHEMISTRY PRELIM 2021 ANS SCHEME

| | | | |
|-----------|----------|-----------|----------|
| 1 | C | 21 | C |
| 2 | B | 22 | B |
| 3 | D | 23 | B |
| 4 | A | 24 | B |
| 5 | A | 25 | D |
| 6 | C | 26 | B |
| 7 | D | 27 | C |
| 8 | D | 28 | A |
| 9 | A | 29 | B |
| 10 | A | 30 | B |
| 11 | A | 31 | D |
| 12 | A | 32 | A |
| 13 | C | 33 | D |
| 14 | B | 34 | C |
| 15 | B | 35 | A |
| 16 | B | 36 | A |
| 17 | B | 37 | C |
| 18 | A | 38 | B |
| 19 | D | 39 | A |
| 20 | D | 40 | C |

6092 Prelim 2021 Answer Key

Paper 1 B

| | | | | | |
|-----|------|-------|-------|-------|-------|
| 1-5 | 6-10 | 11-15 | 16-20 | 21-25 | 26-30 |
|-----|------|-------|-------|-------|-------|

Note to all students:

Markers for national exams have commented on students' handwriting that are too small to be read. Let's have some consideration for the people reading and grading your scripts.

Section A

| Qn | Answer | Marks | Remarks | | | | | | | | | | | | |
|----------------------|---|-------------|---------|------|--------------------|----|-------|----------------------|-------------------|-------------|------------------|-----------------|---------|------------------------------|--|
| A1a | Element X: Group VII / Halogens Element Y: Group VI | 1 1 | | | | | | | | | | | | | |
| b | Oxygen | 1 | | | | | | | | | | | | | |
| c | Period 3 | 1 | | | | | | | | | | | | | |
| d | AlX ₃ | 1 | | | | | | | | | | | | | |
| e | It is inert OR It has low boiling and melting point. | 1 | | | | | | | | | | | | | |
| A2a | <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%;">name</th> <th style="width: 30%;">formula</th> <th style="width: 40%;">code</th> </tr> </thead> <tbody> <tr> <td>Potassium fluoride</td> <td>KF</td> <td>19, 9</td> </tr> <tr> <td>Copper(II) carbonate</td> <td>CuCO₃</td> <td>29, 6, 8(3)</td> </tr> <tr> <td>nitrogen dioxide</td> <td>NO₂</td> <td>7, 8(2)</td> </tr> </tbody> </table> | name | formula | code | Potassium fluoride | KF | 19, 9 | Copper(II) carbonate | CuCO ₃ | 29, 6, 8(3) | nitrogen dioxide | NO ₂ | 7, 8(2) | Every 2 correct answers = 1m | |
| name | formula | code | | | | | | | | | | | | | |
| Potassium fluoride | KF | 19, 9 | | | | | | | | | | | | | |
| Copper(II) carbonate | CuCO ₃ | 29, 6, 8(3) | | | | | | | | | | | | | |
| nitrogen dioxide | NO ₂ | 7, 8(2) | | | | | | | | | | | | | |
| bi | Atoms of the same element (same number of protons) with different number of neutrons. | 1 | | | | | | | | | | | | | |
| bii | There is a higher percentage of Cl-37 atoms compared to Cl-35 atoms in the gas sample. Both Cl-35 and Cl-37 contain 17 protons in the nucleus and 17 electrons in the electron shells. Cl-35 contains 18 neutrons in the nucleus but Cl-37 contains 20 neutrons in the nucleus. | 1 | | | | | | | | | | | | | |
| A3a | CaO(s) + H ₂ O(l) → Ca(OH) ₂ (aq) Correct equation | 1 | | | | | | | | | | | | | |

6092 Prelim 2021 Answer Key

| Qn | Answer | Marks | Remarks |
|-----|--|--------|---------|
| | Correct state symbols | 1 | |
| b | Calcium hydroxide that is formed in a wet concrete beam is a strong alkali that fully dissociates in water to release hydroxide ions. | ½ ½ | |
| c | Carbon dioxide in air is an acidic oxide. It dissolves in water to form a weak acid that will neutralize the calcium hydroxide found in wet concrete. | 1 1 | |
| d | Salt speeds up the process of rusting. | 1 | |
| A4a | $\text{Ag}^+(\text{aq}) + \text{Cl}^-(\text{aq}) \rightarrow \text{AgCl}(\text{s})$ $\text{Ba}^{2+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s})$ | 1 1 | |
| b | Deduct 1m overall if state symbols not provided or wrongly stated. Nitric acid which contains the nitrate ion has been added to the resulting mixture already. The test will always give positive results regardless of whether the original sample of wastewater contained nitrate ions. | 1 1 | |
| C | <p>(1/2m)</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 5px; text-align: center;"> Add <u>excess aqueous ammonia</u> to the sample of wastewater </div> <div style="text-align: center;"> ↑ </div> <div style="border: 1px solid black; padding: 5px; text-align: center;"> A white precipitate is formed which is insoluble in excess aqueous ammonia </div> </div> <p style="text-align: center;">↓ Filter</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 5px; text-align: center;"> White precipitate is formed, insoluble in excess aqueous sodium hydroxide <p>(1/2m)</p> </div> <div style="text-align: center;"> ↓ </div> <div style="border: 1px solid black; padding: 5px; text-align: center;"> Add aqueous sodium hydroxide in <u>excess</u> to the filtrate <p>(1/2m)</p> </div> </div> | 1 | |
| A5a | Production of ammonia in Haber process OR Production of margarine with vegetable oil. | 1 | |
| b | Accept any other reasonable suggestion. X = activation energy | 1 | |

| Qn | Answer | Marks | Remarks | | | | | | | | | | | | | |
|------------|---|--|------------|--|---|------------|--|----------------|---|---|------------|---|----------------|---|------------------|--|
| | Y = enthalpy change | 1 | | | | | | | | | | | | | | |
| c | Agree. The energy profile diagram shows that the energy level of the products are higher than the energy level of the reactants. This means that the reaction is endothermic and there should be greater energy absorbed for breaking bonds than energy released from bond formation. | 1 1 1 | | | | | | | | | | | | | | |
| d | Ea lowered But same start and end point level. | 1 | | | | | | | | | | | | | | |
| A6a | <table border="1"> <thead> <tr> <th>experiment</th> <th>electrodes</th> <th>ionic equation, for reaction at each electrode</th> </tr> </thead> <tbody> <tr> <td rowspan="2">1</td> <td>carbon rod</td> <td>$4\text{OH}^-(\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^-$</td> </tr> <tr> <td>steel keychain</td> <td>$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$</td> </tr> <tr> <td rowspan="2">2</td> <td>copper rod</td> <td>$\text{Cu}(\text{s}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{e}^-$</td> </tr> <tr> <td>steel keychain</td> <td>$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$</td> </tr> </tbody> </table> | experiment | electrodes | ionic equation, for reaction at each electrode | 1 | carbon rod | $4\text{OH}^-(\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^-$ | steel keychain | $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$ | 2 | copper rod | $\text{Cu}(\text{s}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{e}^-$ | steel keychain | $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$ | 1 ½ 1 ½ | |
| experiment | electrodes | ionic equation, for reaction at each electrode | | | | | | | | | | | | | | |
| 1 | carbon rod | $4\text{OH}^-(\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^-$ | | | | | | | | | | | | | | |
| | steel keychain | $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$ | | | | | | | | | | | | | | |
| 2 | copper rod | $\text{Cu}(\text{s}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{e}^-$ | | | | | | | | | | | | | | |
| | steel keychain | $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$ | | | | | | | | | | | | | | |
| b | <p>Experiment 1: The electrolyte solution turns from blue to colourless. This is because, copper(II) ions that cause the blue colouration is being discharged and removed at the cathode.</p> <p>Experiment 2: The electrolyte solution remains blue. This is because, as 1 mole of copper(II) ions are being discharged at the cathode. 1 mole of copper(II) ions are being released/formed at the anode to replenish it.</p> | 1 1 1 1 | | | | | | | | | | | | | | |
| c | The some parts of the original steel material had dissolved away. Steel which contain iron, is more reactive than copper. Iron will displace copper(II) ions out of the solution and form iron(II) ions instead, which dissolves into the solution. | 1 1 1 | | | | | | | | | | | | | | |
| 7a | General formula: $\text{C}_n\text{H}_{2n}\text{O}$ Functional group: C=O | 1 1 | | | | | | | | | | | | | | |

| Qn | Answer | Marks | Remarks |
|----|--|--------|---------|
| b | Approx 104 °C Answer must approximate to the interval difference of the data. | 1 | |
| c | It indicates the position of the carbon where the functional group is found. | 1 | |
| d | 2-propanone and propanal are isomers. They have the same chemical formula of C ₃ H ₆ O. But they have different structures because the oxygen atom is bonded to the 2 nd carbon in 2-propanone but in propanal, the oxygen atom is bonded to the 1 st /3 rd carbon. | 1 1 | |

Section B

| Qn | Answer | Marks | Comments |
|-----|---|--------|----------|
| B8a | Correct sharing of electrons Correct distribution of unshared electrons and correct ratio of elements. | 1 1 | |
| bi | Conc of hydrogen peroxide = 1.0×34 = 34 g/dm ³ | 1 | |
| bii | From ans in bi, 1000 cm ³ of the solution contains 34 g of H ₂ O ₂ . 1000 cm ³ of the solution contains (34/1.45 = 23.448 cm ³) of H ₂ O ₂ . Volume concentration of H ₂ O ₂ = (23.448 / 1000) x 100 = 2.345 % | ½ 1 | |
| c | Since concentration is > 0.5%, it meets the minimum concentration required. Low boiling and melting point Does not conduct electricity | 1 1 | |
| d | For bleach with TEAD: Between 10 – 40 °C, as temperature increases, the effective bleaching performance increases but starts to drop slightly as temperature increases beyond 40 °C. | ½ ½ | |

| Qn | Answer | Marks | Comments |
|-----|---|--------------------|----------|
| | <p>For bleach without TEAD: Between 10 – 35 °C, as temperature increases, the effective bleaching performance decreases but increases beyond 35 °C</p> | <p>1/2 1/2</p> | |
| ei | | 1 | |
| eii | <p>The catalyst increases the proportion of colliding particles with energy equal or more than E_a.</p> | 1 | |
| B9a | <p>On Venus, carbon dioxide is the most abundant gas (96.5%) whereas on earth, carbon dioxide is only a very small 0.03% of the atmosphere.</p> <p>On Venus, nitrogen is a small percentage of 3.5 of the atmosphere but on earth, nitrogen is the most abundant gas (78%).</p> <ul style="list-style-type: none"> - Quote values or difference in gas composition - Provide a description on the significance of the different. (eg. Most abundant etc). | <p>1 1</p> | |
| b | <p>The atmosphere of Venus is mostly made up of carbon dioxide (96.5%) which is a greenhouse gas.</p> | 1 | |

| Qn | Answer | Marks | Comments |
|-----------|--|------------------|----------|
| | Carbon dioxide <u>traps the heat</u> on the planet, causing temperatures to be high. | 1 | |
| c | The material must have a high melting point. This is to withstand the high surface area temperature of Venus and prevent the probe from melting when it lands on the planet. | 1 1 | |
| d | Bubble a sample of the air from Venus through acidified potassium manganate(VII). The purple solution will turn colourless if sulfur dioxide is present. | 1 1 | |
| E B10a | Mass of I ₂ = 127.57 – 126.55 = 1.02 g moles of I ₂ = 1.02 / 254 = 0.0040158 = 0.00402 (3 sf) | 1 1 | |
| b | From chemical formula XI ₂ , mole ratio of X : I ₂ = 1:1 Therefore, no. of moles of X in final dried product = 0.0040158 (same as I ₂) Mass of X in final dried product = 1.28 – 1.02 = 0.26 g 0.0040158 mole of X = 0.26 g 1 mole of X (molar mass) = 0.26 / 0.0040158 = 64.7 = 65 The metal is zinc. | 1 1 1 1 | |
| c | Test the electricity conductivity of the solid compound and the aqueous compound. If the compound is unable to conduct electricity in solid state but able to conduct electricity in aqueous state, the compound is likely to be ionic in nature. | | |
| d | Chlorine is above iodine in Group VII. Since <u>reactivity decreases down the group</u> for halogens, chlorine will be <u>more reactive than iodine</u> and react more vigorously. | 1/2 1 1/2 | |

6092 Prelim 2021 Answer Key

| Qn | Answer | Marks | Comments |
|-------------|--|------------------|----------|
| OR B10ai | $ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{C} = \text{C} \\ \quad \\ \text{H} \quad \text{H} \end{array} $ | 1 | |
| all | $ \begin{array}{c} \text{H} \\ \\ \text{H} - \text{C} - \text{H} \\ \\ \text{O} = \text{C} \\ \quad \\ \text{H} \quad \text{O} \\ \quad \\ \text{C} = \text{C} \\ \quad \\ \text{H} \quad \text{H} \end{array} $ | 1 | |
| b | <p>Total Mr = 12040</p> <p>Mass of poly(vinyl acetate) = $0.2 \times 12040 = 2408$</p> <p>Mr of VA monomer = 86</p> <p>$m = 2408 / 86 = 28$</p> <p>no. of carbon atoms in VA = $28 \times 4 = 112$</p> <p>Mass of poly(ethylene) = $0.8 \times 12040 = 9632$</p> <p>Mr of ethylene monomer = 28</p> <p>$n = 9632 / 28 = 344$</p> <p>no. of carbon atoms in ethylene monomer = $344 \times 2 = 688$</p> <p>Total number of carbon atoms = $688 + 112 = 800$</p> <p>Correct mass found based on %</p> <p>Correct method of finding n and m</p> <p>Correct no. of carbon atoms for the total of each monomer type</p> <p>Correct final answer.</p> | 1 1 1 1 | |

6092 Prelim 2021 Answer Key

| Qn | Answer | Marks | Comments |
|------|--|-------|----------|
| ci | $ \begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H}-\text{C}-\text{O}-\text{H} \\ \\ \text{O} \end{array} $ | 2 | |
| cii | Concentrated sulfuric acid catalyst and warming | 1 | |
| ciii | Bubbles will be formed | 1 | |