



Bukit Batok Secondary School
PRELIMINARY EXAMINATION 2018
SEC 4 EXPRESS

CHEMISTRY

Paper 1 Multiple Choice

6092/01

24 August 2018

0745 - 0845

1 hour

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, glue or correction fluid.

Write your name, index number and class on the Answer Sheet in the spaces provided.

There are **forty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

A copy of the Periodic Table is printed at the end of the question paper.

The use of an approved scientific calculator is expected, where appropriate.

- 1 The reaction scheme shows how hydrated copper(II) sulfate, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, changes when heated.



A little water was accidentally spilled into a dish containing hydrated copper(II) sulfate. What could be done to remove the water, leaving pure, dry $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$?

- A Heat the dish over a boiling water-bath.
 B Heat the dish to a constant mass.
 C Heat the dish with a Bunsen burner.
 D Let the dish stand in direct sunlight.
- 2 Aluminium sulfate is sometimes used in water treatment to remove impurities. Aqueous aluminium sulfate is acidic. The table shows the results of tests on four different samples of treated water.
 To which sample had an excess of aluminium sulfate been added?


sample	pH of sample	reaction with an excess of aqueous ammonia
A	3	white precipitate
B	3	no reaction
C	7	no reaction
D	11	white precipitate

- 3 An acid, X, was added to a solution of the nitrate of a metal, Y. A dense white precipitate was formed.
 What are X and Y?

	acid X	metal Y
A	hydrochloric	calcium
B	nitric	zinc
C	sulfuric	aluminium
D	sulfuric	barium

- 4 A student tested a solution by adding aqueous sodium hydroxide. A precipitate was not seen because the reagent was added too quickly.
 What could **not** have been present in the solution?
 A Al^{3+} B Ca^{2+} C NH_4^+ D Zn^{2+}

- 5 The graph gives the melting points (m.p.) of mixtures of lead and tin.



The graph shows that any mixture of lead and tin must have a melting point that is

- A above that of tin.
 B below that of lead.
 C below that of both tin and lead.
 D between that of tin and lead.
- 6 The isotopes of carbon and oxygen are given in the table.

Isotopes of carbon	^{12}C	^{13}C	^{14}C
Isotopes of oxygen	^{16}O	^{17}O	^{18}O

A molecule of carbon dioxide with molecular mass 46 could contain

- A one ^{12}C atom and two ^{16}O atoms.
 B one ^{14}C atom and two ^{18}O atoms.
 C one ^{12}C atom, one ^{16}O atom and one ^{18}O atom.
 D one ^{14}C atom, one ^{16}O atom and one ^{18}O atom.
- 7 Particles with the same electron arrangement are said to be isoelectronic. Which of the following compounds contains ions which are isoelectronic?
- A CaCl_2
 B KBr
 C MgCl_2
 D Na_2S

- 8 The table shows information about particles X and Y.


	number of protons	number of neutrons	electronic structure
X	9	10	2, 8
Y	17	20	2, 8, 8

Which statement is correct for both X and Y?

- A They are atoms of metals.
 B They are atoms of noble gases.
 C They are isotopes of the same element.
 D They are negative ions.
- 9 The table shows some properties of four substances.
 Which substance is an ionic compound?

	melting point /°C	conducts electricity when solid	dissolves in water	conducts electricity in aqueous solution
A	-102	no	yes	yes
B	801	no	yes	yes
C	842	yes	yes	yes
D	3000	yes	no	no

- 10 The shapes and names of some molecules are shown below.





Phosphine is a compound of phosphorus, an element in Group V, and hydrogen. The shape of a molecule of phosphine is likely to be

- A bent.
 B linear.
 C pyramidal.
 D tetrahedral.

- 11 Which sulfide contains the greatest mass of sulfur in a 10 g sample?

sulfide	formula	mass of one mole /g
A	NiS	91
B	FeS ₂	120
C	MoS ₂	160
D	PbS	239


- 12 Two of the reactions used in the manufacture of nitric acid, HNO₃, are shown.



What is the maximum number of moles of nitric acid which could be formed from one mole of nitrogen monoxide, NO?

- A** 0.5 **B** 1.0 **C** 2.0 **D** 4.0
- 13 A piece of chalk has a mass of 23.0 g. Chalk is impure calcium carbonate. When analysed, the chalk is found to contain 0.226 moles of calcium carbonate. What is the percentage purity of the piece of chalk?
- A** 0.983% **B** 1.02% **C** 77.0% **D** 98.3%
- 14 Which element requires the smallest number of electrons for one mole of atoms to be liberated during electrolysis?
- A** aluminium
B calcium
C copper
D sodium


- 15 The diagrams show an electrolysis experiment using inert electrodes.



Which could be liquid Y?


- 1 aqueous copper(II) sulfate
 - 2 aqueous sodium nitrate
 - 3 concentrated aqueous sodium chloride
 - 4 dilute sulfuric acid
- A** 4 only
B 1 and 4 only
C 2 and 4 only
D 2, 3 and 4 only
- 16 A student carries out a single experiment to determine the speed of reaction between calcium carbonate and an excess of hydrochloric acid. Which of the following does **not** change during the course of the reaction?
- A** concentration of the hydrochloric acid solution
 - B** mass of the calcium carbonate
 - C** volume of carbon dioxide evolved
 - D** volume of hydrochloric acid solution

- 17 The graph shows how the rate of a reaction varies with the concentration of one of the reactants.



What is the reaction time, in seconds, when the concentration of the reactant was 0.50 mol/dm³?

- A 0.2 B 0.5 C 2.0 D 5.0
- 18 A diagram for the energy change during a chemical reaction is shown.



For which reaction(s) would this be an appropriate diagram?

- 1 $\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$
- 2 $\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O}$
- 3 $2\text{C} + \text{O}_2 \rightarrow 2\text{CO}$

- A 1 only
 B 1 and 2 only
 C 1 and 3 only
 D 1, 2 and 3

19 The oxide of titanium, TiO_2 , is used as a 'whitener' in toothpaste. It is obtained from the ore iron(II) titanate, FeTiO_3 . What is the change, if any, in the oxidation number of titanium in the reaction $\text{FeTiO}_3 \rightarrow \text{TiO}_2$?

- A It is oxidized from +3 to +4.
- B It is reduced from +3 to +2.
- C It is reduced from +6 to +4.
- D There is no change in the oxidation number.

20 The pH of an aqueous solution of hydrochloric acid is 2. What will be the pH of the acid after the addition of 10.0 g of sodium chloride?

- A 2 B 5 C 7 D 9

21 Which row in the table correctly shows the properties of 0.100 mol/dm^3 hydrochloric acid when compared with 0.100 mol/dm^3 ethanoic acid?

	pH	conductivity	Rate of reaction with magnesium
A	lower	lower	slower
B	higher	higher	faster
C	lower	higher	faster
D	higher	lower	slower

22 Consider the three reactions below.

- reaction between nitric acid and calcium hydroxide
- ethane burning in air
- reaction between ethanoic acid and ethanol


A student made three statements about the three reactions above.

- 1 carbon dioxide is produced in all reactions
- 2 water is produced in all reactions
- 3 a salt is produced in all reactions

Which statement(s) is/are true?

- A 2 only
- B 1 and 2 only
- C 2 and 3 only
- D 1, 2 and 3

- 23 Methylamine, CH_3NH_2 , has very similar chemical properties to ammonia, NH_3 . Methylamine reacts with hydrogen chloride to form a white crystalline salt, methylammonium chloride.




A sample of methylammonium chloride is heated with aqueous sodium hydroxide. What are the products?

- A ammonia, sodium chloride and water
 B ammonia, sodium hydrogencarbonate and sodium chloride
 C methylamine, hydrogen chloride and water
 D methylamine, sodium chloride and water
- 24 A student has five reagents.
- dilute hydrochloric acid
 - dilute sulfuric acid
 - dilute nitric acid
 - solid calcium carbonate
 - solid copper(II) carbonate
- How many soluble salts can be prepared?
- A 3 B 4 C 5 D 6
- 25 How can a pure sample of barium sulfate be obtained from barium carbonate?
- A Dissolve it in dilute hydrochloric acid, add dilute sulfuric acid, filter and crystallise.
 B Dissolve it in dilute hydrochloric acid, add dilute sulfuric acid, filter and wash.
 C Dissolve it in water, add dilute sulfuric acid, filter and crystallise.
 D Dissolve it in water, add dilute sulfuric acid, filter and wash.

- 26 An alloy of copper and zinc is added to an excess of dilute hydrochloric acid. The resulting mixture is then filtered. Which observations are correct?

	filtrate	residue
A	colourless solution	none
B	colourless solution	pinkish brown
C	blue solution	grey
D	blue solution	none

- 27 In the apparatus shown, gas P is passed over solid Q.



No reaction occurs if P and Q are

	P	Q
A	hydrogen	lead(II) oxide
B	hydrogen	magnesium oxide
C	oxygen	carbon
D	oxygen	sulfur

- 28 The period 4 elements gallium (Ga), germanium (Ge), arsenic (As) and selenium (Se) are elements below aluminium, silicon, phosphorus and sulfur in the Periodic Table, a portion of which is shown below.

period 3 elements	Al	Si	P	S
period 4 elements	Ga	Ge	As	Se

The properties of each period 4 element resemble those of the period 3 element directly above it.


Which period 4 elements form oxides that dissolve in water to give an acid solution?

- A As and Se
 B Ga and Ge
 C Ga and Se
 D Se only
- 29 When a mineral was heated in a Bunsen flame to a constant mass, a colourless gas that produced a white precipitate in limewater, was given off. The remaining solid was cooled and then added to aqueous hydrochloric acid. Vigorous effervescence was seen. What was the mineral?
- A aragonite, CaCO_3
 B artinite, $\text{MgCO}_3 \cdot \text{Mg}(\text{OH})_2 \cdot 3\text{H}_2\text{O}$
 C barytocalcite, $\text{BaCO}_3 \cdot \text{CaCO}_3$
 D dolomite, $\text{CaCO}_3 \cdot \text{MgCO}_3$

- 30 Listed below are four solutions.
- 1 aqueous sodium hydroxide
 - 2 aqueous silver nitrate
 - 3 aqueous potassium sulfate
 - 4 dilute hydrochloric acid

Which of the following solution(s) will react with magnesium metal?

- A 4 only
 B 1 and 4
 C 2 and 4 only
 D 2, 3 and 4 only
- 31 Attaching pieces of magnesium to underground iron pipes can protect the iron from corrosion.
 Which reaction protects the iron from corrosion?
- A $\text{Fe}^{2+}(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + \text{e}$
 B $\text{Fe}(\text{s}) \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}$
 C $\text{Mg}^{2+}(\text{aq}) + 2\text{e} \rightarrow \text{Mg}(\text{s})$
 D $\text{Mg}(\text{s}) \rightarrow \text{Mg}^{2+}(\text{aq}) + 2\text{e}$
- 32 An experiment to find the percentage of oxygen in 150 cm^3 of polluted air is shown.



The apparatus is left for one week.

After this time, the volume of gas in the measuring cylinder is 122 cm^3 .

What is the percentage of oxygen, to the nearest whole number, in the polluted air?

- A 19% B 21% C 28% D 81%
- 33 The depletion of the ozone layer in the upper atmosphere reduces the Earth's natural protection from harmful ultraviolet radiation.
 Which compound would cause the most depletion of the ozone layer?
- A CCl_3F B CF_4 C CHClF_2 D CH_2F_2

- 34 The compound, C_8H_{18} undergoes the following process.




Which row in the table correctly identifies Process X and Compound Y?

	Process X	Compound Y
A	cracking	hexane
B	cracking	hexene
C	distillation	hexane
D	distillation	hexene

- 35 How many moles of hydrogen chloride are formed when one mole of methane is added to a large excess of chlorine in the dark?

A 0 **B** 1 **C** 2 **D** 4

- 36 Three members of the cycloalkene homologous series are shown:



Which of the following is the general formula for this homologous series?


- A** C_nH_{2n-4}
B C_nH_{2n-2}
C C_nH_{2n}
D C_nH_{2n+2}

- 37 Oil contains carbon-carbon double bonds which can undergo addition reactions with iodine. The iodine number of an oil is the mass of iodine in grams that will react with 100 g of oil.

Which row in the table shows the oil that is likely to have the lowest melting point?

	oil	iodine number
A	corn	123
B	linseed	179
C	olive	81
D	soya	130

- 38 A compound has the following structure.



Which reaction(s) will occur with this compound?

- 1 Bromine water will decolourise.
- 2 It will react with an alcohol to form an ester.
- 3 It will react with sodium metal.

- A** 1 only
B 1 and 2 only
C 2 and 3 only
D 1, 2 and 3


- 39 Polyvinyl chloride (PVC) is a man-made polymer used mainly in the manufacture of pipes. PVC pipes are strong, lightweight and does not rot.

Which statements correctly describe the polymer, polyvinyl chloride, PVC?

- 1 Combustion of PVC waste produces a highly acidic gas.
- 2 PVC molecules are saturated.
- 3 The empirical formula of PVC is the same as the empirical formula of its monomers.

- A** 1 and 2 only
B 1 and 3 only
C 2 and 3 only
D 1, 2 and 3





40. A section of a condensation polymer is shown below.



One of the monomers is



The structural formula of the other monomer is



End of paper

The Periodic Table of Elements

		Group																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
I	II	III	IV	V	VI	VII	0					0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
3 Li lithium 7	4 Be beryllium 9	11 Na sodium 23	12 Mg magnesium 24	13 Al aluminium 27	14 Si silicon 28	15 P phosphorus 31	16 S sulfur 32	17 Cl chlorine 35.5	18 Ar argon 40	19 K potassium 39	20 Ca calcium 40	21 Sc scandium 45	22 Ti titanium 48	23 V vanadium 51	24 Cr chromium 52	25 Mn manganese 55	26 Fe iron 56	27 Co cobalt 59	28 Ni nickel 59	29 Cu copper 64	30 Zn zinc 65	31 Ga gallium 70	32 Ge germanium 73	33 As arsenic 75	34 Se selenium 79	35 Br bromine 80	36 Kr krypton 84	37 Rb rubidium 85	38 Sr strontium 88	39 Y yttrium 89	40 Zr zirconium 91	41 Nb niobium 93	42 Mo molybdenum 96	43 Tc technetium -	44 Ru ruthenium 101	45 Rh rhodium 103	46 Pd palladium 106	47 Ag silver 108	48 Cd cadmium 112	49 In indium 115	50 Sn tin 119	51 Sb antimony 122	52 Te tellurium 128	53 I iodine 127	54 Xe xenon 131	55 Cs caesium 133	56 Ba barium 137	57-71 lanthanoids	72 Hf hafnium 178	73 Ta tantalum 181	74 W tungsten 184	75 Re rhenium 186	76 Os osmium 190	77 Ir iridium 192	78 Pt platinum 195	79 Au gold 197	80 Hg mercury 201	81 Tl thallium 204	82 Pb lead 207	83 Bi bismuth 209	84 Po polonium -	85 At astatine -	86 Rn radon -	87 Fr francium -	88 Ra radium -	89-103 actinoids	104 Rf rutherfordium -	105 Db dubnium -	106 Sg seaborgium -	107 Bh bohrium -	108 Hs hassium -	109 Mt meitnerium -	110 Ds darmstadtium -	111 Rg roentgenium -	112 Cn copernicium -	116 Lv livermorium -	117 Ts tennessine -	118 Og oganeson -	119 Nh nihonium -	120 Fl flerovium -	121 Mc moscovium -	122 Lv livermorium -	123 Ts tennessine -	124 Og oganeson -	125 Nh nihonium -	126 Fl flerovium -	127 Mc moscovium -	128 Lv livermorium -	129 Ts tennessine -	130 Og oganeson -	131 Nh nihonium -	132 Fl flerovium -	133 Mc moscovium -	134 Lv livermorium -	135 Ts tennessine -	136 Og oganeson -	137 Nh nihonium -	138 Fl flerovium -	139 Mc moscovium -	140 Lv livermorium -	141 Ts tennessine -	142 Og oganeson -	143 Nh nihonium -	144 Fl flerovium -	145 Mc moscovium -	146 Lv livermorium -	147 Ts tennessine -	148 Og oganeson -	149 Nh nihonium -	150 Fl flerovium -	151 Mc moscovium -	152 Lv livermorium -	153 Ts tennessine -	154 Og oganeson -	155 Nh nihonium -	156 Fl flerovium -	157 Mc moscovium -	158 Lv livermorium -	159 Ts tennessine -	160 Og oganeson -	161 Nh nihonium -	162 Fl flerovium -	163 Mc moscovium -	164 Lv livermorium -	165 Ts tennessine -	166 Og oganeson -	167 Nh nihonium -	168 Fl flerovium -	169 Mc moscovium -	170 Lv livermorium -	171 Ts tennessine -	172 Og oganeson -	173 Nh nihonium -	174 Fl flerovium -	175 Mc moscovium -	176 Lv livermorium -	177 Ts tennessine -	178 Og oganeson -	179 Nh nihonium -	180 Fl flerovium -	181 Mc moscovium -	182 Lv livermorium -	183 Ts tennessine -	184 Og oganeson -	185 Nh nihonium -	186 Fl flerovium -	187 Mc moscovium -	188 Lv livermorium -	189 Ts tennessine -	190 Og oganeson -	191 Nh nihonium -	192 Fl flerovium -	193 Mc moscovium -	194 Lv livermorium -	195 Ts tennessine -	196 Og oganeson -	197 Nh nihonium -	198 Fl flerovium -	199 Mc moscovium -	200 Lv livermorium -	201 Ts tennessine -	202 Og oganeson -	203 Nh nihonium -	204 Fl flerovium -	205 Mc moscovium -	206 Lv livermorium -	207 Ts tennessine -	208 Og oganeson -	209 Nh nihonium -	210 Fl flerovium -	211 Mc moscovium -	212 Lv livermorium -	213 Ts tennessine -	214 Og oganeson -	215 Nh nihonium -	216 Fl flerovium -	217 Mc moscovium -	218 Lv livermorium -	219 Ts tennessine -	220 Og oganeson -	221 Nh nihonium -	222 Fl flerovium -	223 Mc moscovium -	224 Lv livermorium -	225 Ts tennessine -	226 Og oganeson -	227 Nh nihonium -	228 Fl flerovium -	229 Mc moscovium -	230 Lv livermorium -	231 Ts tennessine -	232 Og oganeson -	233 Nh nihonium -	234 Fl flerovium -	235 Mc moscovium -	236 Lv livermorium -	237 Ts tennessine -	238 Og oganeson -	239 Nh nihonium -	240 Fl flerovium -	241 Mc moscovium -	242 Lv livermorium -	243 Ts tennessine -	244 Og oganeson -	245 Nh nihonium -	246 Fl flerovium -	247 Mc moscovium -	248 Lv livermorium -	249 Ts tennessine -	250 Og oganeson -	251 Nh nihonium -	252 Fl flerovium -	253 Mc moscovium -	254 Lv livermorium -	255 Ts tennessine -	256 Og oganeson -	257 Nh nihonium -	258 Fl flerovium -	259 Mc moscovium -	260 Lv livermorium -	261 Ts tennessine -	262 Og oganeson -	263 Nh nihonium -	264 Fl flerovium -	265 Mc moscovium -	266 Lv livermorium -	267 Ts tennessine -	268 Og oganeson -	269 Nh nihonium -	270 Fl flerovium -	271 Mc moscovium -	272 Lv livermorium -	273 Ts tennessine -	274 Og oganeson -	275 Nh nihonium -	276 Fl flerovium -	277 Mc moscovium -	278 Lv livermorium -	279 Ts tennessine -	280 Og oganeson -	281 Nh nihonium -	282 Fl flerovium -	283 Mc moscovium -	284 Lv livermorium -	285 Ts tennessine -	286 Og oganeson -	287 Nh nihonium -	288 Fl flerovium -	289 Mc moscovium -	290 Lv livermorium -	291 Ts tennessine -	292 Og oganeson -	293 Nh nihonium -	294 Fl flerovium -	295 Mc moscovium -	296 Lv livermorium -	297 Ts tennessine -	298 Og oganeson -	299 Nh nihonium -	300 Fl flerovium -	301 Mc moscovium -	302 Lv livermorium -	303 Ts tennessine -	304 Og oganeson -	305 Nh nihonium -	306 Fl flerovium -	307 Mc moscovium -	308 Lv livermorium -	309 Ts tennessine -	310 Og oganeson -	311 Nh nihonium -	312 Fl flerovium -	313 Mc moscovium -	314 Lv livermorium -	315 Ts tennessine -	316 Og oganeson -	317 Nh nihonium -	318 Fl flerovium -	319 Mc moscovium -	320 Lv livermorium -	321 Ts tennessine -	322 Og oganeson -	323 Nh nihonium -	324 Fl flerovium -	325 Mc moscovium -	326 Lv livermorium -	327 Ts tennessine -	328 Og oganeson -	329 Nh nihonium -	330 Fl flerovium -	331 Mc moscovium -	332 Lv livermorium -	333 Ts tennessine -	334 Og oganeson -	335 Nh nihonium -	336 Fl flerovium -	337 Mc moscovium -	338 Lv livermorium -	339 Ts tennessine -	340 Og oganeson -	341 Nh nihonium -	342 Fl flerovium -	343 Mc moscovium -	344 Lv livermorium -	345 Ts tennessine -	346 Og oganeson -	347 Nh nihonium -	348 Fl flerovium -	349 Mc moscovium -	350 Lv livermorium -	351 Ts tennessine -	352 Og oganeson -	353 Nh nihonium -	354 Fl flerovium -	355 Mc moscovium -	356 Lv livermorium -	357 Ts tennessine -	358 Og oganeson -	359 Nh nihonium -	360 Fl flerovium -	361 Mc moscovium -	362 Lv livermorium -	363 Ts tennessine -	364 Og oganeson -	365 Nh nihonium -	366 Fl flerovium -	367 Mc moscovium -	368 Lv livermorium -	369 Ts tennessine -	370 Og oganeson -	371 Nh nihonium -	372 Fl flerovium -	373 Mc moscovium -	374 Lv livermorium -	375 Ts tennessine -	376 Og oganeson -	377 Nh nihonium -	378 Fl flerovium -	379 Mc moscovium -	380 Lv livermorium -	381 Ts tennessine -	382 Og oganeson -	383 Nh nihonium -	384 Fl flerovium -	385 Mc moscovium -	386 Lv livermorium -	387 Ts tennessine -	388 Og oganeson -	389 Nh nihonium -	390 Fl flerovium -	391 Mc moscovium -	392 Lv livermorium -	393 Ts tennessine -	394 Og oganeson -	395 Nh nihonium -	396 Fl flerovium -	397 Mc moscovium -	398 Lv livermorium -	399 Ts tennessine -	400 Og oganeson -	401 Nh nihonium -	402 Fl flerovium -	403 Mc moscovium -	404 Lv livermorium -	405 Ts tennessine -	406 Og oganeson -	407 Nh nihonium -	408 Fl flerovium -	409 Mc moscovium -	410 Lv livermorium -	411 Ts tennessine -	412 Og oganeson -	413 Nh nihonium -	414 Fl flerovium -	415 Mc moscovium -	416 Lv livermorium -	417 Ts tennessine -	418 Og oganeson -	419 Nh nihonium -	420 Fl flerovium -	421 Mc moscovium -	422 Lv livermorium -	423 Ts tennessine -	424 Og oganeson -	425 Nh nihonium -	426 Fl flerovium -	427 Mc moscovium -	428 Lv livermorium -	429 Ts tennessine -	430 Og oganeson -	431 Nh nihonium -	432 Fl flerovium -	433 Mc moscovium -	434 Lv livermorium -	435 Ts tennessine -	436 Og oganeson -	437 Nh nihonium -	438 Fl flerovium -	439 Mc moscovium -	440 Lv livermorium -	441 Ts tennessine -	442 Og oganeson -	443 Nh nihonium -	444 Fl flerovium -	445 Mc moscovium -	446 Lv livermorium -	447 Ts tennessine -	448 Og oganeson -	449 Nh nihonium -	450 Fl flerovium -	451 Mc moscovium -	452 Lv livermorium -	453 Ts tennessine -	454 Og oganeson -	455 Nh nihonium -	456 Fl flerovium -	457 Mc moscovium -	458 Lv livermorium -	459 Ts tennessine -	460 Og oganeson -	461 Nh nihonium -	462 Fl flerovium -	463 Mc moscovium -	464 Lv livermorium -	465 Ts tennessine -	466 Og oganeson -	467 Nh nihonium -	468 Fl flerovium -	469 Mc moscovium -	470 Lv livermorium -	471 Ts tennessine -	472 Og oganeson -	473 Nh nihonium -	474 Fl flerovium -	475 Mc moscovium -	476 Lv livermorium -	477 Ts tennessine -	478 Og oganeson -	479 Nh nihonium -	480 Fl flerovium -	481 Mc moscovium -	482 Lv livermorium -	483 Ts tennessine -	484 Og oganeson -	485 Nh nihonium -	486 Fl flerovium -	487 Mc moscovium -	488 Lv livermorium -	489 Ts tennessine -	490 Og oganeson -	491 Nh nihonium -	492 Fl flerovium -	493 Mc moscovium -	494 Lv livermorium -	495 Ts tennessine -	496 Og oganeson -	497 Nh nihonium -	498 Fl flerovium -	499 Mc moscovium -	500 Lv livermorium -	501 Ts tennessine -	502 Og oganeson -	503 Nh nihonium -	504 Fl flerovium -	505 Mc moscovium -	506 Lv livermorium -	507 Ts tennessine -	508 Og oganeson -	509 Nh nihonium -	510 Fl flerovium -	511 Mc moscovium -	512 Lv livermorium -	513 Ts tennessine -	514 Og oganeson -	515 Nh nihonium -	516 Fl flerovium -	517 Mc moscovium -	518 Lv livermorium -	519 Ts tennessine -	520 Og oganeson -	521 Nh nihonium -	522 Fl flerovium -	523 Mc moscovium -	524 Lv livermorium -	525 Ts tennessine -	526 Og oganeson -	527 Nh nihonium -	528 Fl flerovium -	529 Mc moscovium -	530 Lv livermorium -	531 Ts tennessine -	532 Og oganeson -	533 Nh nihonium -	534 Fl flerovium -	535 Mc moscovium -	536 Lv livermorium -	537 Ts tennessine -	538 Og oganeson -	539 Nh nihonium -	540 Fl flerovium -	541 Mc moscovium -	542 Lv livermorium -	543 Ts tennessine -	544 Og oganeson -	545 Nh nihonium -	546 Fl flerovium -	547 Mc moscovium -	548 Lv livermorium -	549 Ts tennessine -	550 Og oganeson -	551 Nh nihonium -	552 Fl flerovium -	553 Mc moscovium -	554 Lv livermorium -	555 Ts tennessine -	556 Og oganeson -	557 Nh nihonium -	558 Fl flerovium -	559 Mc moscovium -	560 Lv livermorium -	561 Ts tennessine -	562 Og oganeson -	563 Nh nihonium -	564 Fl flerovium -	565 Mc moscovium -	566 Lv livermorium -	567 Ts tennessine -	568 Og oganeson -	569 Nh nihonium -	570 Fl flerovium -	571 Mc moscovium -	572 Lv livermorium -	573 Ts tennessine -	574 Og oganeson -	575 Nh nihonium -	576 Fl flerovium -	577 Mc moscovium -	578 Lv livermorium -	579 Ts tennessine -	580 Og oganeson -	581 Nh nihonium -	582 Fl flerovium -	583 Mc moscovium -	584 Lv livermorium -	585 Ts tennessine -	586 Og oganeson -	587 Nh nihonium -	588 Fl flerovium -	589 Mc moscovium -	590 Lv livermorium -	591 Ts tennessine -	592 Og oganeson -	593 Nh nihonium -	594 Fl flerovium -	595 Mc moscovium -	596 Lv livermorium -	597 Ts tennessine -	598 Og oganeson -	599 Nh nihonium -	600 Fl flerovium -	601 Mc moscovium -	602 Lv livermorium -	603 Ts tennessine -	604 Og oganeson -	605 Nh nihonium -	606 Fl flerovium -	607 Mc moscovium -	608 Lv livermorium -	609 Ts tennessine -	610 Og oganeson -	611 Nh nihonium -	612 Fl flerovium -	613 Mc moscovium -	614 Lv livermorium -	615 Ts tennessine -	616 Og oganeson -	617 Nh nihonium -	618 Fl flerovium -	619 Mc moscovium -	620 Lv livermorium -	621 Ts tennessine -	622 Og oganeson -	623 Nh nihonium -	624 Fl flerovium -	625 Mc moscovium -	626 Lv livermorium -	627 Ts tennessine -	628 Og oganeson -	629 Nh nihonium -	630 Fl flerovium -	631 Mc moscovium -	632 Lv livermorium -	633 Ts tennessine -	634 Og oganeson -	635 Nh nihonium -	636 Fl flerovium -	637 Mc moscovium -	638 Lv livermorium -	639 Ts tennessine -	640 Og oganeson -	641 Nh nihonium -	642 Fl flerovium -	643 Mc moscovium -	644 Lv livermorium -	645 Ts tennessine -	646 Og oganeson -	647 Nh nihonium -	648 Fl flerovium -	649 Mc moscovium -	650 Lv livermorium -	651 Ts tennessine -	652 Og oganeson -	653 Nh nihonium -	654 Fl flerovium -	655 Mc moscovium -	656 Lv livermorium -	657 Ts tennessine -	658 Og oganeson -	659 Nh nihonium -	660 Fl flerovium -	661 Mc moscovium

Name: Index No. Class:



Bukit Batok Secondary School
PRELIMINARY EXAMINATION 2018
Sec 4 EXPRESS

CHEMISTRY

Paper 2

6092/02**15 August 2018****1030 - 1215****1 hour 45 minutes**

Candidates answer on the Question Paper.
 No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your name, index number and class in the spaces provided at the top of this page.

Write in dark blue or black pen

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Section A

Answer **all** questions in the spaces provided.

Section B

Answer **all three** questions, the last question is in the form of either/or.

Answer **all** questions in the spaces provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

A copy of the Periodic Table is given at the end of the paper.

The use of an approved scientific calculator is expected, where appropriate.

For Examiner's use	
Section A	/50
Section B	
B7	
B8	
B9	
Total	/80

This document consists of 23 printed pages.

Section A

Answer all the questions in this section in the spaces provided.

The total mark for this section is 50

A1(a) The grid below represents part of a blank periodic table, the numbers being the proton number of the elements.

In the grid below, write

- (i) **P** in a space which could be occupied by a noble gas which is used to fill weather balloons. [1]
- (ii) **Q** in a space which the most reactive non-metal would occupy. [1]
- (iii) **R** in a space which could be occupied by a metal with the lowest density. [1]
- (iv) **S** in a space which could be occupied by an element forming an amphoteric hydroxide. [1]
- (v) **T** in a space which could be occupied by an element with an isotope that can be represented by $^{14}_6X$. [1]

1							2
3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18

- (b) (i) Describe how the metallic character of the elements in Period 3 changes across the period from left to right.

.....[1]

- (ii) State how the metallic character of an element is related to its electronic structure.

..... [1]

- (c) Explain what is meant by the term *periodicity*.

.....

.....[1]

[Total: 8]

A2 Carbon atoms can bond to each other to produce a variety of different structures, including diamond, graphite and buckminsterfullerene.

(a) There are similarities and differences in the structure and bonding in diamond and graphite.

(i) Describe two features of the structure and bonding in diamond that are similar to graphite.


.....
.....
.....
..... [2]

(ii) Describe two features of the structure and bonding in diamond that are different from graphite.

.....
.....
.....
..... [2]

(b) Buckminsterfullerene is a form carbon with the formula C_{60} . If it is burned completely in oxygen, it forms carbon dioxide as the only product. Calculate the mass of carbon dioxide that is released when 51 g of buckminsterfullerene is completely burned in oxygen. [2]

(c) The structures of graphite and boron nitride are shown below.



- (i) What is the chemical formula for boron nitride?[1]
- (ii) Like graphite, boron nitride feels slippery to the touch.
Explain, in terms of bonding and structure, why boron nitride feels slippery to touch.


.....

.....

.....

..... [2]

- (iii) The diagram below shows the structure of a solid form of carbon called graphene. Graphene contains **one layer** of carbon atoms. Graphene is made from graphite but it is harder than graphite.



Explain, using ideas about structure and bonding, why graphene is hard.

.....

..... [1]


[Total: 10]

A3 Alkanes like propane and butane are found in Liquefied Petroleum Gases(LPG).

- (a) An experiment shows that complete combustion of 1.0 dm^3 (measured at room temperature and pressure) of butane produces 120 kJ of energy. Calculate a value for the enthalpy change of complete combustion (kJ/mol) of butane, giving the correct sign.

[1]

- (b)(i) The alkane with 5 carbon atoms, pentane exists as several isomers shown below. One is straight chain pentane while the other two are branched chain pentane.



Will the two isomers which are branched chain pentane have the same enthalpy change on complete combustion as the straight chain pentane?

Explain your reasoning.

.....
 [1]

- (ii) The table shows the enthalpy changes of combustion of hexane and heptane.

name	formula	enthalpy change of combustion / kJ/mol
hexane	C_6H_{14}	-4163
heptane	C_7H_{16}	-4817

Using the data given, estimate the enthalpy change of combustion in kJ/mol of octane, C_8H_{18} . Explain the method you use to arrive at your answer.

.....

 [2]

- (c) Some students studied the graph below that shows the amount of fossil fuel burned in the world between 1960 and 2010.




Fig 3.1

- (i) One student says that the amount of fossil fuels burned has increased by the same amount every ten years.
Is the student correct? Use data from the graph to justify your answer.

.....
 [1]

- (ii) Another student says that it is very difficult to estimate the amount of fossil fuel we will use in 100 years' time. Suggest reasons the student could give to justify this statement.

.....

 [2]

- (iii) The graph below shows the changes in average global temperature from 1960 to 2010.




Fig 3.2

Describe the link between the trends shown in the graphs in Fig 3.1 and Fig 3.2.

.....


.....[1]

[Total: 8]

A4 The diagram shows a simple cell, with two different metals as electrodes dipped in dilute nitric acid. A student did an experiment using the simple cell below.

The voltages were recorded in the table.

- If the voltage measured is positive then metal 2 is more reactive than metal 1.
- If the voltage measured is negative then metal 1 is more reactive than metal 2.



		metal 2				
		beryllium	cobalt	nickel	silver	vanadium
metal 1	beryllium	0.0 V	-1.6 V	-1.6 V	not measured	-0.7 V
	cobalt		0.0 V	0.0 V	-1.1 V	0.9 V
	nickel			0.0 V	-1.1 V	0.9 V
	silver				0.0 V	2.0 V
	vanadium					0.0 V

(a) (i) In the simple cell containing nickel and silver, it was observed that the electrolyte slowly turned pale green. Write the ionic equation to explain the colour change.

.....[1]

(ii) What happened to the mass of the nickel electrode?

.....[1]

(b) (i) Using the data given, state the most reactive metal in the table above. Explain your reasoning.

.....

.....

.....[2]

(ii) Predict the voltage produced by a simple cell with beryllium as metal 1 and silver as metal 2.

.....[1]

- (c) (i) The student wanted to rank the metals listed in the table according to their reactivity but he was not able to do so. Why?

.....[1]


- (ii) Briefly describe one **simple** experiment the student can do which will help him to solve the problem in c(i).

.....

.....[1]

[Total: 7]

- A5 The rate of reaction of iron with aqueous bromine is determined by using the apparatus shown below.



The iron is removed at regular intervals. It is washed, dried and then weighed. The iron is then replaced in the solution.

The experiment is repeated twice, each time with a different concentration of aqueous bromine at room temperature, 25 °C. The results are shown in the table below.

Experiment	concentration of aqueous bromine mol/dm ³	speed of reaction mg iron reacted/min
1	0.050	9.2
2	0.10	18.1
3	0.15	27.2

- (a) Describe how and explain why the speed of this reaction changes with concentration of bromine.

.....

.....

.....[2]

(b) (i) Experiment 1 is repeated after aqueous bromine has been cooled in an ice bath to 15°C.
Predict the speed of reaction, with appropriate unit.....[1]

(ii) Using collision theory, explain your answer in (b)(i).

.....

.....

.....

..... [2]


(c) Suggest another method for measuring the speed of this reaction.

.....[1]

[Total: 6]

A6 Ammonia, NH₃, is a colourless, pungent-smelling gas which has been known to man from the beginning of recorded time. Chemists have discovered a novel way of 'fixing' atmospheric nitrogen (converting nitrogen gas into its compounds). Moist nitrogen is passed over a TiO₂ plate which has been coated with other chemicals. The nitrogen is thought to react with moisture in the air at room temperature and pressure to form ammonia.

A possible equation for the reaction is given below.



(a) (i) Explain why there are only a few reactions that 'fix' nitrogen.

.....

.....

..... [2]

(ii) Suggest and explain one advantage and one disadvantage of the process given in the equation above as a method of making ammonia compared with the Haber process.

.....

.....

.....

..... [2]

(b) 1.20 dm³ of ammonia gas was dissolved in water to form 200 cm³ of aqueous alkali at room temperature and pressure.


(i) Calculate how many moles of NH₃(g) was dissolved in water.

[1]

(ii) Write the equation for the neutralisation of aqueous ammonia, NH₃(aq) by dilute sulfuric acid.

..... [1]

(c) The decomposition of ammonia is represented by the following equation.




The activation energy, E_a for the uncatalysed reaction is 335 kJ/mol.

The activation energy, E_a for the reaction when tungsten is used as a catalyst is 163 kJ/mol.

(i) On the grid provided on page 12, draw a **labelled** energy profile diagram for the uncatalysed and catalysed reactions.

[3]


Include the necessary information given.



- (ii) When osmium is used as a catalyst, the activation energy is 197 kJ/mol. Which catalyst, osmium or tungsten, will cause ammonia to decompose at a faster rate? Explain your answer using ideas about particles.

.....

- (iii) State the activation energy for the uncatalysed reaction of the following: [1]



..... [1]
 [Total:11]

Section B

Answer all three questions in this section.

The last question is in the form of an either/or and only one of the alternatives should be attempted.

B7 Composition of sea water

The Earth's ocean holds about 1.5×10^{18} tonnes of water, which in turn contains 0.05×10^{18} tonnes of dissolved salts. The table below shows eight most common ions in the sea.

ion	% by mass of total dissolved solids	concentration in mol/dm ³
Chloride, Cl ⁻	55.04	0.535
Sodium, Na ⁺	30.42	0.457
Sulfate, SO ₄ ²⁻	7.69	0.028
Magnesium, Mg ²⁺	3.91	0.056
Calcium, Ca ²⁺	1.16	0.010
Potassium, K ⁺	1.10	0.0097
Carbonate, CO ₃ ²⁻	0.41	0.0023
Bromide, Br ⁻	0.19	0.00081

The dissolved ions in the sea form an essentially free source of materials to anyone with access to the sea. Evaporation of sea water produces sodium chloride and potassium chloride. The two other elements that can be obtained from sea water are bromine and magnesium.


Uses of Magnesium

Magnesium is the lightest structural metal used today, some 30% lighter than aluminium. Magnesium is the third most used metal in construction (after iron and aluminium). Nearly 70% of the world production of magnesium is used to make alloys. One example is Magnox which is an alloy of magnesium with small amount of aluminium and other metals.

Extraction of magnesium


The first stage in the production of magnesium is to mix the sea water with a slurry of calcium hydroxide. This precipitates magnesium hydroxide.

This reaction can be represented as follows.




Magnesium hydroxide is then separated and heated to produce magnesium oxide.

The flow chart summarises the process mentioned



Conversion to magnesium chloride is achieved by heating the oxide, mixed with carbon, in a stream of chlorine at a high temperature in the furnace.



The resulting anhydrous magnesium chloride is fed into electrolytic cells. A schematic diagram of the electrolytic cell is shown below in Fig 7.1




Fig 7.1

The design of this cell considers the following properties of both magnesium metal and magnesium chloride:

- molten magnesium reacts vigorously with oxygen
- at the temperature of molten magnesium chloride, magnesium is a liquid
- molten magnesium has a lower density than molten magnesium chloride and forms a separate layer on the surface.

- (a) Name the most abundant ionic compound in sea water and determine the effective concentration of this compound in mol/dm³.

.....
 [1]

- (b) (i) From the information given, deduce the trend in solubility of the Group II metal hydroxide as the proton number increases.

.....
 [1]

- (ii) Calculate the mass of magnesium hydroxide precipitated when an excess of calcium hydroxide is added to 1000 dm³ of sea water.

[2]

- (c) (i) Write an ionic equation for the reaction at the cathode in the electrolytic cell.

..... [1]

- (ii) How does the design of the cell shown in the Fig 7.1, take into consideration the reaction of molten magnesium with oxygen?

.....
 [1]

- (d) Electrolysis is an expensive process as high consumption of energy is needed. Using the information given, what is being done to lower the cost in industrial process?

.....
 [1]

B8 (a) A student reacted together an alcohol and a carboxylic acid under appropriate conditions to produce an ester.
A sweet smelling organic liquid, **Q**, with the empirical formula C_2H_4O was produced.
The M_r of **Q** was found by experiments to be 87.5.

(i) What is the molecular formula of **Q**? Show the necessary calculation. [1]

(ii) In the boxes below, draw the structural formula of **two** isomers with this formula that are **straight chain** esters. [2]

--	--

A sample of **Q** was heated with aqueous sulfuric acid. The product obtained was a mixture of the original alcohol and carboxylic acid. This mixture was heated under reflux with acidified potassium manganate(VII) to give a **single** product, **R**.

The product, **R**, was collected and subjected to the following tests:

- A sample of **R** gave no reaction with aqueous bromine.
- A second sample of **R** gave an effervescence with sodium carbonate.
- A third sample of **R** is completely miscible with water.


(iii) What is the identity of single organic compound **R**?

..... [1]

(b) Cyclopentene is a cyclic alkene with the formula C_5H_8 . It is a colourless liquid with a petrol-like odour. It is used as a monomers for synthesis of plastics.

The figure below shows some reactions involving cyclopentene

(i) Complete the partial structures of compounds X and Y which are the products of the reactions. [2]



(ii) Write a balanced chemical equation to show the reaction between cyclopentene and aqueous bromine.

..... [1]

(iii) Cyclopentene can be polymerised to give poly(cyclopentene). Draw a section of poly(cyclopentene) to show two repeat units. [1]

[Total: 8]

B9 Either

Aqueous iron(II) bromide is a pale green solution containing iron(II) ions and bromide ions. When chlorine is passed into aqueous iron(II) bromide, the colour of the solution changes from pale green to orange-red.

When the orange-red solution is heated, it gives off a brown vapour, leaving a yellow solution **S**. The brown vapour forms a dark orange liquid **T** on cooling. When ethene gas is bubbled into **T**, the dark orange colour disappears. Sodium hydroxide solution is added to solution **S** and a reddish brown precipitate was obtained.

- (a) (i) Name liquid **T**. [1]
 (ii) Draw 'dot-and-cross' diagram to show the electron arrangement in **T**.
 Show only the outer electrons. [1]

- (b) Name the yellow compound present in solution **S**.
 [1]

- (c) (i) Construct a balanced chemical equation for the reaction in which **S** and **T** are formed.
 [1]


- (ii) In this reaction in which **S** and **T** are formed, name the oxidising agent.
 Explain your answer, using **electron transfer**.

.....

 [2]

- (d) A student uses chromatography to analyse the food dyes used in a packet of sweets. The packet contains three different coloured sweets. He tests one sweet of each colour. He uses two known safe food dyes and one known unsafe dye as references.

The chromatogram below shows his results.



- (i) The student looks at the results and makes this statement:
 "The results show that it is possible that two of the sweets contain an unsafe dye."
 Explain how the results of the chromatogram support the student's conclusion.

.....

 [2]

- (ii) Calculate the R_f value of the unsafe dye given in the chromatogram above.

- (iii) The student also uses chromatography to identify the **flavourings** used in the sweets. He sprays his chromatogram with a locating agent.
 Why does he need to use a locating agent?

..... [1]

B9 Or

Both calcium and barium are elements in Group II of the Periodic Table. The trend of the reactivity of the elements in Group II is similar to that in Group I. Like Group I elements, calcium and barium form salts with the halogens.

The salt, calcium chloride, CaCl_2 , can be made by different reactions.

A student prepared hydrated calcium chloride by carrying out the following experiment.

- Step 1** The student added an excess of a solid calcium compound, **X**, to dilute hydrochloric acid. The mixture fizzed as the solid reacted.
- Step 2** The student filtered the mixture to give an aqueous solution of CaCl_2 .
- Step 3** On evaporation, colourless crystals of hydrated calcium chloride were formed.

(a) Why is calcium chloride an example of 'salt'?

.....
..... [1]

(b) A friend of the student suggested that solid **X** was calcium oxide. State one reason why the student's friend was **incorrect** and suggest a possible identity of solid **X**.

.....
.....
..... [2]

(c) Hydrated calcium chloride has a molar mass of 219 g/mol.
Determine the formula of **hydrated** calcium chloride.
You must show your working.

[2]

- (d) Calcium chloride can also be formed by directly reacting calcium with chlorine gas. Explain, using **oxidation states**, why the formation of calcium chloride from its elements, is a redox reaction.

.....
.....
.....[2]

- (e) The student decided to prepare barium sulfate, BaSO_4 , by adding barium metal to dilute sulfuric acid. Another student said this method should not be used to prepare the salt, barium sulfate. Give **two** reasons why the other student is correct.

.....
.....
.....[2]

- (f) Barium atom has the electron arrangement 2, 8, 18, 18, 8, 2. Write the electron arrangement of the barium **ion**.

.....[1]

End of paper

Group																																																																																																																																																																																																																																																																																																																																																																																							
I	II	III	IV	V	VI	VII	0																																																																																																																																																																																																																																																																																																																																																																																
3 Li lithium 7	4 Be beryllium 9	11 Na sodium 23	12 Mg magnesium 24	19 K potassium 39	20 Ca calcium 40	37 Rb rubidium 85	38 Sr strontium 88	55 Cs caesium 133	87 Fr francium -	21 Sc scandium 45	39 Y yttrium 89	57-71 lanthanoids	88-103 actinoids	1 H hydrogen 1	5 B boron 11	6 C carbon 12	7 N nitrogen 14	8 O oxygen 16	9 F fluorine 19	10 Ne neon 20	13 Al aluminium 27	14 Si silicon 28	15 P phosphorus 31	16 S sulfur 32	17 Cl chlorine 35.5	18 Ar argon 40	25 Mn manganese 55	26 Fe iron 56	27 Co cobalt 59	28 Ni nickel 59	29 Cu copper 64	30 Zn zinc 65	41 Nb niobium 93	42 Mo molybdenum 96	43 Tc technetium -	44 Ru ruthenium 101	45 Rh rhodium 103	46 Pd palladium 106	47 Ag silver 108	48 Cd cadmium 112	49 In indium 115	50 Sn tin 119	51 Sb antimony 122	52 Te tellurium 128	53 I iodine 127	54 Xe xenon 131	56 Ba barium 137	72 Hf hafnium 178	73 Ta tantalum 181	74 W tungsten 184	75 Re rhenium 186	76 Os osmium 190	77 Ir iridium 192	78 Pt platinum 195	79 Au gold 197	80 Hg mercury 201	81 Tl thallium 204	82 Pb lead 207	83 Bi bismuth 209	84 Po polonium -	85 At astatine -	86 Rn radon -	89-103 actinoids	89 Ac actinium -	90 Th thorium 232	91 Pa protactinium 231	92 U uranium 238	93 Np neptunium -	94 Pu plutonium -	95 Am americium -	96 Cm curium -	97 Bk berkelium -	98 Cf californium -	99 Es einsteinium -	100 Fm fermium -	101 Md mendelevium -	102 No nobelium -	103 Lr lawrencium -	104 Rf rutherfordium -	105 Db dubnium -	106 Sg seaborgium -	107 Bh bohrium -	108 Hs hassium -	109 Mt meitnerium -	110 Ds darmstadtium -	111 Rg roentgenium -	112 Cn copernicium -	113 Nh nihonium -	114 Fl flerovium -	115 Lv livermorium -	116 Ts tennessium -	117 Og oganesson -	118 Uue unbinilium -	119 Uuh ununilium -	120 Uuq ununquadium -	121 Uub ununbium -	122 Uut ununtrium -	123 Uuq ununquadium -	124 Uub ununbium -	125 Uut ununtrium -	126 Uuq ununquadium -	127 Uub ununbium -	128 Uut ununtrium -	129 Uuq ununquadium -	130 Uub ununbium -	131 Uut ununtrium -	132 Uuq ununquadium -	133 Uub ununbium -	134 Uut ununtrium -	135 Uuq ununquadium -	136 Uub ununbium -	137 Uut ununtrium -	138 Uuq ununquadium -	139 Uub ununbium -	140 Uut ununtrium -	141 Uuq ununquadium -	142 Uub ununbium -	143 Uut ununtrium -	144 Uuq ununquadium -	145 Uub ununbium -	146 Uut ununtrium -	147 Uuq ununquadium -	148 Uub ununbium -	149 Uut ununtrium -	150 Uuq ununquadium -	151 Uub ununbium -	152 Uut ununtrium -	153 Uuq ununquadium -	154 Uub ununbium -	155 Uut ununtrium -	156 Uuq ununquadium -	157 Uub ununbium -	158 Uut ununtrium -	159 Uuq ununquadium -	160 Uub ununbium -	161 Uut ununtrium -	162 Uuq ununquadium -	163 Uub ununbium -	164 Uut ununtrium -	165 Uuq ununquadium -	166 Uub ununbium -	167 Uut ununtrium -	168 Uuq ununquadium -	169 Uub ununbium -	170 Uut ununtrium -	171 Uuq ununquadium -	172 Uub ununbium -	173 Uut ununtrium -	174 Uuq ununquadium -	175 Uub ununbium -	176 Uut ununtrium -	177 Uuq ununquadium -	178 Uub ununbium -	179 Uut ununtrium -	180 Uuq ununquadium -	181 Uub ununbium -	182 Uut ununtrium -	183 Uuq ununquadium -	184 Uub ununbium -	185 Uut ununtrium -	186 Uuq ununquadium -	187 Uub ununbium -	188 Uut ununtrium -	189 Uuq ununquadium -	190 Uub ununbium -	191 Uut ununtrium -	192 Uuq ununquadium -	193 Uub ununbium -	194 Uut ununtrium -	195 Uuq ununquadium -	196 Uub ununbium -	197 Uut ununtrium -	198 Uuq ununquadium -	199 Uub ununbium -	200 Uut ununtrium -	201 Uuq ununquadium -	202 Uub ununbium -	203 Uut ununtrium -	204 Uuq ununquadium -	205 Uub ununbium -	206 Uut ununtrium -	207 Uuq ununquadium -	208 Uub ununbium -	209 Uut ununtrium -	210 Uuq ununquadium -	211 Uub ununbium -	212 Uut ununtrium -	213 Uuq ununquadium -	214 Uub ununbium -	215 Uut ununtrium -	216 Uuq ununquadium -	217 Uub ununbium -	218 Uut ununtrium -	219 Uuq ununquadium -	220 Uub ununbium -	221 Uut ununtrium -	222 Uuq ununquadium -	223 Uub ununbium -	224 Uut ununtrium -	225 Uuq ununquadium -	226 Uub ununbium -	227 Uut ununtrium -	228 Uuq ununquadium -	229 Uub ununbium -	230 Uut ununtrium -	231 Uuq ununquadium -	232 Uub ununbium -	233 Uut ununtrium -	234 Uuq ununquadium -	235 Uub ununbium -	236 Uut ununtrium -	237 Uuq ununquadium -	238 Uub ununbium -	239 Uut ununtrium -	240 Uuq ununquadium -	241 Uub ununbium -	242 Uut ununtrium -	243 Uuq ununquadium -	244 Uub ununbium -	245 Uut ununtrium -	246 Uuq ununquadium -	247 Uub ununbium -	248 Uut ununtrium -	249 Uuq ununquadium -	250 Uub ununbium -	251 Uut ununtrium -	252 Uuq ununquadium -	253 Uub ununbium -	254 Uut ununtrium -	255 Uuq ununquadium -	256 Uub ununbium -	257 Uut ununtrium -	258 Uuq ununquadium -	259 Uub ununbium -	260 Uut ununtrium -	261 Uuq ununquadium -	262 Uub ununbium -	263 Uut ununtrium -	264 Uuq ununquadium -	265 Uub ununbium -	266 Uut ununtrium -	267 Uuq ununquadium -	268 Uub ununbium -	269 Uut ununtrium -	270 Uuq ununquadium -	271 Uub ununbium -	272 Uut ununtrium -	273 Uuq ununquadium -	274 Uub ununbium -	275 Uut ununtrium -	276 Uuq ununquadium -	277 Uub ununbium -	278 Uut ununtrium -	279 Uuq ununquadium -	280 Uub ununbium -	281 Uut ununtrium -	282 Uuq ununquadium -	283 Uub ununbium -	284 Uut ununtrium -	285 Uuq ununquadium -	286 Uub ununbium -	287 Uut ununtrium -	288 Uuq ununquadium -	289 Uub ununbium -	290 Uut ununtrium -	291 Uuq ununquadium -	292 Uub ununbium -	293 Uut ununtrium -	294 Uuq ununquadium -	295 Uub ununbium -	296 Uut ununtrium -	297 Uuq ununquadium -	298 Uub ununbium -	299 Uut ununtrium -	300 Uuq ununquadium -	301 Uub ununbium -	302 Uut ununtrium -	303 Uuq ununquadium -	304 Uub ununbium -	305 Uut ununtrium -	306 Uuq ununquadium -	307 Uub ununbium -	308 Uut ununtrium -	309 Uuq ununquadium -	310 Uub ununbium -	311 Uut ununtrium -	312 Uuq ununquadium -	313 Uub ununbium -	314 Uut ununtrium -	315 Uuq ununquadium -	316 Uub ununbium -	317 Uut ununtrium -	318 Uuq ununquadium -	319 Uub ununbium -	320 Uut ununtrium -	321 Uuq ununquadium -	322 Uub ununbium -	323 Uut ununtrium -	324 Uuq ununquadium -	325 Uub ununbium -	326 Uut ununtrium -	327 Uuq ununquadium -	328 Uub ununbium -	329 Uut ununtrium -	330 Uuq ununquadium -	331 Uub ununbium -	332 Uut ununtrium -	333 Uuq ununquadium -	334 Uub ununbium -	335 Uut ununtrium -	336 Uuq ununquadium -	337 Uub ununbium -	338 Uut ununtrium -	339 Uuq ununquadium -	340 Uub ununbium -	341 Uut ununtrium -	342 Uuq ununquadium -	343 Uub ununbium -	344 Uut ununtrium -	345 Uuq ununquadium -	346 Uub ununbium -	347 Uut ununtrium -	348 Uuq ununquadium -	349 Uub ununbium -	350 Uut ununtrium -	351 Uuq ununquadium -	352 Uub ununbium -	353 Uut ununtrium -	354 Uuq ununquadium -	355 Uub ununbium -	356 Uut ununtrium -	357 Uuq ununquadium -	358 Uub ununbium -	359 Uut ununtrium -	360 Uuq ununquadium -	361 Uub ununbium -	362 Uut ununtrium -	363 Uuq ununquadium -	364 Uub ununbium -	365 Uut ununtrium -	366 Uuq ununquadium -	367 Uub ununbium -	368 Uut ununtrium -	369 Uuq ununquadium -	370 Uub ununbium -	371 Uut ununtrium -	372 Uuq ununquadium -	373 Uub ununbium -	374 Uut ununtrium -	375 Uuq ununquadium -	376 Uub ununbium -	377 Uut ununtrium -	378 Uuq ununquadium -	379 Uub ununbium -	380 Uut ununtrium -	381 Uuq ununquadium -	382 Uub ununbium -	383 Uut ununtrium -	384 Uuq ununquadium -	385 Uub ununbium -	386 Uut ununtrium -	387 Uuq ununquadium -	388 Uub ununbium -	389 Uut ununtrium -	390 Uuq ununquadium -	391 Uub ununbium -	392 Uut ununtrium -	393 Uuq ununquadium -	394 Uub ununbium -	395 Uut ununtrium -	396 Uuq ununquadium -	397 Uub ununbium -	398 Uut ununtrium -	399 Uuq ununquadium -	400 Uub ununbium -

Key
 proton (atomic) number
 atomic symbol
 name
 relative atomic mass

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

2018 GCE O Prelim sec 4E Chemistry 6092 Paper 1**Answer & mark scheme**

No.	Ans	Remarks
1	D	(A) A boiling water bath has a temperature of 100°C, so if heat over water bath, CuSO ₄ .H ₂ O obtained (see eqn) (B) This will convert hydrated copper(II) sulfate to the anhydrous form. (C) Same as (B)
2	A	(See QA notes)
3	D	X has sulfate ion reacts with barium ion to form insoluble barium sulfate. This is not the reaction of the hydrogen ion in the acid.
4	B	Both aluminium ion and zinc ion forms white ppt which dissolves in excess aq NaOH giving colourless solution. Calcium ion forms white ppt that does not dissolve in excess aq NaOH.
5	B	
6	C	(add up nucleon no. given in the options and compare to 46)
7	D	(write down electron arrangement of ions given in the option and compare) Electron arrangement for: Ca ²⁺ 2,8,8 Cl ⁻ 2,8,8
8	D	(compare no. of protons and no. of electrons) X: 9 protons, 10 electrons so X is negative ion Y: 17 protons, 18 electrons, so Y is negative ion
9	B	(see the column for ability to conduct electricity in solid and in aq state)
10	C	(compare with N which is in Group V, compound of N and H is NH ₃ , so compound of P and H is PH ₃ . Check the no. of white dots which represent hydrogen atoms)
11	B	Mass of S = $\frac{\text{no. of S atom} \times \text{Ar of S}}{\text{Mr}}$ x mass of sample
12	B	
13	D	Mass of calcium carbonate in the chalk = 0.226 x 100 = 22.6g % purity = 22.6 / 23.0 x 100 = 98.3%
14	D	(compare the charge of the positive ion, eg Na ⁺ + e → Na)
15	C	Oxygen and hydrogen gas given off (see volumes of gas produced), so hydroxide ion and hydrogen ion discharged. In (1), copper(II) ion discharged instead of hydrogen ion. In (3), chloride ion discharged instead of hydroxide ion due to higher [Cl ⁻]
16	D	As reaction progresses: (A) Concentration of acid drops (B) More carbonate used (C) More gas produced
17	D	Reaction time, t = 1 / rate = 1 / 0.2 = 5.0 s
18	D	(2) is respiration which is exothermic. (see glucose react with oxygen. Both (1) and (3) are combustion of fuel so exothermic.

No.	Ans	Remarks
19	D	(assign oxidation numbers) In FeTiO_3 , oxidation number of Ti is +4, in TiO_2 , oxidation number is +4
20	A	Sodium chloride is neutral, does not react with hydrogen ions in the acid, so pH unchanged
21	C	Hydrochloric acid is strong acid, total ionization. ethanoic acid is weak acid, partial ionization.
22	A	Carbon dioxide is only produced in 2 nd reaction. Salt is produced in 1 st reaction.
23	D	(Recall properties of ammonium compound, apply this to methyl ammonium chloride) when ammonium chloride react with aq NaOH, ammonia gas, salt (sodium chloride) and water produced.
24	C	With calcium carbonate, soluble salts will be calcium chloride, calcium nitrate. NOT calcium sulfate as it is insoluble With copper(II) carbonate, soluble salts will be copper(II) chloride, copper(II) sulfate, copper(II) nitrate
25	B	Barium sulfate is insoluble, so need two soluble starting reagents. Barium carbonate is also insoluble. Add barium carbonate to dilute hydrochloric acid to form soluble barium chloride, before reacting with the second soluble reagent.
26	B	Copper will not react, remain as residue. Zinc react with dil hydrochloric acid to form colourless solution zinc chloride.
27	B	(A) Lead(II) oxide will be reduced by hydrogen gas to form lead and water (B) Magnesium is higher up in the reactivity series, so hydrogen is not able to reduce magnesium oxide (C) Carbon react with oxygen to form carbon dioxide (D) Sulfur react with oxygen to form sulfur dioxide
28	A	Both P and S are non-metals, so form acidic oxides which dissolves in water to form an acid.
29	C	Going down group II, the carbonate becomes more difficult to decompose. Both magnesium carbonate and calcium carbonate decomposes, but not barium carbonate.
30	C	(2) magnesium will displace silver from silver nitrate (4) magnesium will react with acid to form salt and hydrogen
31	D	Magnesium is a more reactive metal, so loses electron more easily.
32	A	Rusting uses up oxygen. Volume of oxygen used = $150 - 122 = 28 \text{ cm}^3$ % of oxygen = $28 / 150 \times 100 = 18.7\%$
33	A	Chlorine atom reacts with the ozone molecules, so choose the option with largest no. of chlorine atoms.
34	A	Y is C_6H_{14} , so it is hexane
35	A	Absence of uv light, so no substitution occurs
36	B	Cyclobutene – C_4H_6 , cyclopentane – C_5H_8 , cyclohexene – C_6H_{10}
37	B	The oil with the lowest bp will have the largest no. of C=C bonds.

No.	Ans	Remarks
38	D	It has C=C so will react with aq bromine With the -COOH, it is an organic acid, so react with metal and alcohol
39	D	(1) True, Hydrogen chloride gas produced (2) True, no more C=C in addition polymers (3) True, only the polymer is produced
40	B	

Sec 4E GCE O Prelim Chemistry 6092

Answers & mark scheme

The paper was

A1a(i) P – box 2; Q – box 9; R – box 3; S – box 13 T – box 6 [1] each

(b) (i) elements becomes less metallic. [1]

(ii) metallic elements have fewer outer / valence electrons. [1]

(c) periodicity is a repeating pattern (across different periods) [1]

[Total: 8]

A2a(i) Both consists entirely of **carbon** atoms joined by **covalent** bonds; [1]

Both have giant lattice (or giant molecular) [1]

ii. In diamond every carbon atom is bonded to four other carbon atoms, but in graphite, each carbon atom is bonded to 3 atoms;

diamond has a tetrahedral arrangement of atoms but graphite has alayered arrangement;

graphite has delocalised electrons unlike diamond which do not [any 2]

b. No. of mole of $C_{60} = 51 / 720 = 0.0708$ (eqn – optional)

No. of mole of $CO_2 = 60 \times \text{no. of mole of } C_{60} = 0.0708 \times 60 = 4.24$ [1]

Mass of $C_{60} = 4.24 \times 44 = .187$ g [1]

c(i) BN [1]

ii. weak Van der Waal (or intermolecular) forces of attraction between layers; [1]

layers of **atoms** can slide over each other. [1]

iii. graphene has **many** strong covalent bond between carbon atoms. [1]

[Total: 10]

A3a. $\Delta H = -24 \times 120 = -2880 \text{ kJ/mol}$ [reject if no unit and sign] [1]

b(i) Yes, same(not similar) type bond and same number of bond [1]

ii. difference in $\Delta H = 4817 - 4163 = 654$ kJ [1]

from hexane to heptane, increase in one CH_2 group

from heptane to octane, same increase of one CH_2 group

so ΔH for octane = $-(4187 + 654) = -5471$ kJ/mol [1]

c(i) No, quote any two data that shows a difference for every ten years. [1]

Egs of data that can be used: 1960 -70, 2000 millions of tons bigger than 1970 to 80 which has increase 1200 millions of tons, or 1990 – 2000, increase of 800 millions of tons smaller than 2000 – 2010 increase of 1300 millions of tons.

ii. alternative / renewable forms of energy being used; [1]

fossil fuel running out. [1]

iii. As the amount of fossil fuel burnt increase, the increase average global temperature is higher. [1]

[Total: 8]


A4a(i) $Ni(s) \rightarrow Ni^{2+}(aq) + 2e$ [1]

ii. mass decrease [1]

- b(i) beryllium; [1]
It has the largest voltage with cobalt/nickel [1]
- ii. $-2.7\text{ V (V + Ag) + (V + Be)}$ [1]
- c(i). both nickel and cobalt has the same reactivity [1]
- ii. Place a piece of nickel in cobalt nitrate solution. If nickel displaces cobalt, nickel is more reactive than cobalt. [1]
- [Total: 7]

- A5a. As concentration increases, the speed of this reaction increases. When concentration increases, there is greater number of particles in the same volume [1]
Particles are closer to each other so frequency of effective collision increases. [1]
- b(i) 4.5 – 5.0 mg iron reacted/min (units needed)
- ii. As temperature drops, particles loses energy, move slower. [1]
Number of particles with energy equal to or greater than activation energy drops. [1]
Frequency of effective collision decreases.
- c. measure the colour intensity of aqueous bromine. [1]
- [Total: 6]

- A6a(i) $\text{N}\equiv\text{N}$ triple bond; [1]
A lot of energy is needed to break the (strong covalent) bond [1]
- ii. Advantage: lower temperature / lower pressure so save energy, lesser fossil fuel, or water, instead of hydrogen, water is used, so cheaper [1]
Disadvantage: nitrogen oxide produced, reacts with oxygen to form nitrogen dioxide which contribute to acid rain / an air pollutant [1]
- b(i) no. of moles of ammonia = $1.20 / 24 = 0.05$ [1]
- (ii) $2\text{NH}_3(\text{aq}) + \text{H}_2\text{SO}_4(\text{aq}) \rightarrow (\text{NH}_4)_2\text{SO}_4(\text{aq})$ (state symbols not needed) [1]
[1]
- c(i) correct shape and location for both graphs [1m each]
correct labels of E_a , E_a'' and ΔH [1]



- b(ii) Tungsten: with it, the reaction has a **lower activation energy**;
which means the **higher** proportion of collisions that are successful between ammonia
molecules will be higher (not more collisions, both points needed) [1]
- (iii) $E_a = 243 \text{ kJ/mol}$ (units needed) [1]
- [Total: 11]

Section B

- B7a. sodium chloride, $\text{Conc}(\text{mol}/\text{dm}^3) = 0.457$ (sodium ion is the limiting reactant) [1]
- b(i) as the proton number increases, the group II metal hydroxide becomes
more soluble [1]
- ii. no. of mol of Mg^{2+} in 1000 dm^3 sea water = $0.056 \times 1000 = 56$ moles [1]
no. of mol of $\text{Mg}(\text{OH})_2 = \text{no. of mol of } \text{Mg}^{2+} = 56 \text{ mol}$
mass of $\text{Mg}(\text{OH})_2 = 58 \times 56 = 3248 \text{ g}$ [1]
- c(i) $\text{Mg}^{2+}(\text{l}) + 2\text{e} \rightarrow \text{Mg}(\text{l})$ [1]
- ii an inert gas, instead of air, is blown through the cathode compartment above molten
magnesium [1]
- d. sodium chloride and calcium chloride is added to molten magnesium chloride to **lower
the melting point**, saving energy. [1]
- e. at the anode, silver will be oxidised instead of chloride ion. Silver ion would be
produced rather than chlorine at the anode. [1]
 $\text{Ag}(\text{s}) \rightarrow \text{Ag}^+(\text{l}) + \text{e}$ [1]
the silver ion would move to the cathode, get discharged and silver is produced instead
at the cathode. [1]

- f. chlorine gas produced during electrolysis is used to convert magnesium oxide to magnesium chloride at the furnace. [1]
- g. magnesium ion, aluminium ion, delocalised electron [all three correct -1] [Total: 12]



B8a(i) relative mass of $C_2H_4O = 44$

$M_r \sim 88$


$N = 88 / 44 = 2$

Relative molecular formula is $C_4H_8O_2$ [1]

ii. ethyl ethanoate




Methylpropanoate




propyl methanoate [any 2]

iii. ethanoic acid [1]

b(i)



Compound X




compound Y

[1m for each, all H atoms needed]

ii. $C_5H_8 + Br_2 \rightarrow C_5H_8Br_2$ [1]


iii.



[Must have at least two repeat units and the free bonds at the end. All carbon-carbon bonds in the polymer chain must be shown.] [1]

[Total: 8]

B9 Either

- a(i) bromine [1]
 ii (dot-cross diagram of bromine molecule,  [1]
- b. iron(III) chloride [1]
 c(i) $3\text{Cl}_2 + 2\text{FeBr}_2 \rightarrow 2\text{FeCl}_3 + 2\text{Br}_2$
 [1]
 ii. chlorine [1]
 chlorine removes electrons from iron(II) ion and bromide ion. [1]
- d(i) sweet 2 contains an unsafe dye; [1]
 unknown dye in sweet 3 does not match up with a safe dye [1]
 ii. $0.8 / 5.2 = 0.154$ (or 0.15) [1]
 iii. to see the spots / make the colourless spots visible [1]
 (ignore 'find / identify the spots')

[Total: 10]

B9 Or

- a. Hydrogen ion / H^+ ion in acid replaced by calcium ion / Ca^{2+} ion or metal ion. [1]
 b. The reaction produced a gas / calcium oxide does not produce a gas in reaction with acid. [1]
 calcium carbonate. [1]
 c. Mr of $\text{CaCl}_2 = 111$.

$$\text{No. of water molecules} = \frac{219 - 111}{18}$$
 [1]

$$= 6$$

 Formula: $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ [6 and $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ score the 2nd mark, allow no dot, $\text{CaCl}_2 6\text{H}_2\text{O}$] [1]
 d. Calcium is oxidised as oxidation state of calcium increases from 0 to +2 [1]
 Chlorine is reduced as oxidation state of chlorine decreases from 0 to -1 [1]
 e. barium is very reactive metal, so react violently with the acid, reaction not safe; [1]
 barium sulfate formed is insoluble, so form a barrier on barium, preventing further reaction. [1]
 (ii) 2, 8, 18, 18, 8 [1]

[Total: 10]